
Comparing invariants of $\mathbf{SK}_1$

Abstract

In this text, we compare several invariants of the reduced Whitehead group $\mathbf{SK}_1$ of a central simple algebra.

For biquaternion algebras, we compare a generalised invariant of Suslin as constructed by the author in [Wou] to an invariant introduced by Knus-Merkurjev-Rost-Tignol [KMRT]. Using explicit computations, we prove these invariants are essentially the same.

We also prove the non-triviality of an invariant introduced by Kahn [Kah2]. To obtain this result, we compare Kahn's invariant to an invariant introduced by Suslin in 1991 [Sus1] which is non-trivial for Platonov's examples of non-trivial SK_1 [Pla]. We also give a formula for the value on the centre of the tensor product of two symbol algebras which generalises a formula of Merkurjev for biquaternion algebras [Mer1].

1 Introduction

Let k be a field and A a central simple k -algebra. The triviality of the *reduced Whitehead group* $\mathbf{SK}_1(A)$ (which is isomorphic to $\mathbf{SL}_1(A)/[A^\times, A^\times]$) is a long studied question. Tannaka and Artin posed the question in the 1930's [NM, Wan]. For more than 30 years, one tried to prove the triviality of $\mathbf{SK}_1(A)$ in full generality. In 1976, Platonov gave a counterexample using discrete valuation rings [Pla, Thm. 5.19]. Wang, however, did prove the triviality of $\mathbf{SK}_1(A)$ if $\text{ind}_k(A)$ is square-free [Wan]. This inspired Suslin to conjecture that this can be the only case of triviality [Sus1]. This would give a sufficient answer to the question of Tannaka and Artin. Merkurjev proved it is true when $4 \mid \text{ind}_k(A)$ [Mer3]; and Rehmann-Tikhonov-Yanchevskiĭ proved it is sufficient to prove the conjecture for the tensor product of two symbol algebras [RTY, Thm. 0.19].

In order to study his conjecture, Suslin conjectured in 1991 the existence of a cohomological invariant of $\mathbf{SK}_1(A)$ with values in Galois cohomology ($n = \text{ind}_k(A) \in k^\times$):

$$\rho : \mathbf{SK}_1(A)(k) \rightarrow H^4(k, \mu_n^{\otimes 3}) / (H^2(k, \mu_n^{\otimes 2}) \cup [A]), \quad (1.1)$$

where $[A]$ stands for the class of A in ${}_n\text{Br}(k) \cong H^2(k, \mu_n)$ [Sus1, Conj. 11.6]. There are various definitions of invariants of this flavour. In 1991, Suslin defined twice his conjectured invariant (ibid., §2). For biquaternion algebras Rost gave a closely related invariant [Mer1, Thm. 4], and in 2006 Suslin defined his conjectured invariant in full

K.U.Leuven – Departement Wiskunde, Celestijnenlaan 200B bus 2400, B-3001 Leuven, Belgium – tim@wouters.in

2010 *Mathematics Subject Classification*: 19B99 (12G05, 16K50, 17B20)

Keywords: Reduced Whitehead Group – Suslin's conjecture – Cohomological invariants – Symbol algebras

generality [Sus2, §6]. Kahn even generalised this invariant to a range of new invariants [Kah2, Cor. 8.4 & Def. 11.3].

The restriction to central simple algebras with $n = \text{ind}_k(A) \in k^\times$ is a natural one, since otherwise the cohomology groups can be trivial and ${}_n\text{Br}(k)$ does not have to be isomorphic to $H^2(k, \mu_n)$. Using Kato's cohomology of *logarithmic differentials* [Kat1], the author generalised any of the aforementioned invariants to all central simple algebras using a lift from positive characteristic to characteristic 0 [Wou]. We recall the definitions of the invariants in more detail in Section 2. It is generally assumed that all defined invariants are essentially the same, but very few results exist on this subject. In this paper, we compare some of them.

For biquaternion algebras, Knus-Merkurjev-Rost-Tignol constructed a cohomological invariant of $\mathbf{SK}_1(A)$ without the condition on the index [KMRT, §17]. They use Witt groups, Witt rings, and an involution on the biquaternion algebra to define it. If $\text{char}(k) \neq 2$, they prove the invariant is essentially the same as Suslin's invariant for biquaternions. Using the construction of the generalisation of Suslin's invariant, we prove that for base fields of characteristic 2 their invariant essentially equals Suslin's generalised invariant (Section 3).

In Section 4, we compare a new invariant of Kahn with all of the other existing invariants (or more correctly, we compare the invariants to Kahn's invariant). This allows us to prove the non-triviality of Kahn's invariant for Platonov's examples of non-trivial SK_1 . We also prove a formula for the value on the centre of the tensor product of two symbol algebras under Kahn's invariant which generalises a formula of Merkurjev for biquaternion algebras ([Mer1, Ex. p.70] – see also [KMRT, Ex. 17.23]).

Notations – Let us fix the following notations throughout this text.

- If k is a field, then k_s denotes a separable closure and $\Gamma_k = \text{Gal}(k_s/k)$ its absolute Galois group. Furthermore, denote $\mathbb{G}_m = \text{Spec}(\mathbb{Z}[t, t^{-1}])$.
- A prime factorisation $p_1^{e_1} \cdot \dots \cdot p_r^{e_r}$ of a (positive) integer m is always supposed to be primitive (i.e. $m = p_1^{e_1} \cdot \dots \cdot p_r^{e_r}$, with p_i primes, $e_i \geq 1$ integers for $1 \leq i \leq r$ and $p_i \neq p_j$ for any $1 \leq i < j \leq r$).
- We use standard notations for the following categories: the category **Sets** of sets, the category $k\text{-fields}$ of field extensions of a field k , the category **Groups** of groups, and the category **Ab** of abelian groups.
- If $m > 0$ is an integer (prime to $\text{char}(k)$), then μ_m denotes the Γ_k -module of consisting of m -th roots of unity of k_s . If we want to stress the field in use, we write $\mu_m(k)$ (so that this can be viewed as the k -rational points of the appropriate sheaf).
- The appearing cohomology groups are Galois cohomology groups (unless mentioned otherwise).
- ${}_m\text{Br}(k)$ is the m -th torsion part of the Brauer group of k ($m > 0$ an integer). If K is a field extension of k , we denote by $\text{Br}(K/k)$ the kernel of the base extension morphism $\text{Br}(k) \rightarrow \text{Br}(K)$.

- If F is a discrete valuation field (with valuation v), then the valuation ring is denoted by $\mathcal{O}_v$ and the residue field by $\kappa(v)$. If $x \in \mathcal{O}_v$, we denote by $\bar{x}$ its class in $\kappa(v)$. We also use this notation for other objects for which we can define (canonical) residues. A discrete valuation is supposed to be non-trivial (of rank 1). By F_{nr} we denote the maximal unramified extension of F .
- If A is a central simple k -algebra and if F is a field extension of k , then $A_F = A \otimes_k F$ is the central simple F -algebra obtained from A by base extension to F . More generally, for a ring R , a commutative R -algebra S , and an Azumaya R -algebra A , we denote $A_S = A \otimes_R S$, the Azumaya S -algebra obtained from A by base extension to S . By $[A]$ we denote the Brauer class of a central simple algebra/Azumaya algebra A .
- For any central simple algebra A and $a \in A$, the reduced norm of a is denoted as $\text{Nrd}_{A/k}(a)$. In the same way, $\text{Trd}_{A/k}(a)$ is the reduced trace and $\text{Prd}_{A,a/k}(X)$ is the reduced characteristic polynomial. If

$$\text{Prd}_{A,a/k}(X) = X^n - s_1(a)X^{n-1} + \dots + (-1)^n s_n(a),$$

then we know $\text{Nrd}_{A/k}(a) = s_n(a)$ and $\text{Trd}_{A/k}(a) = s_1(a)$.

- For a central simple k -algebra, we denote by $\mathbf{SL}_1(A)$ the usual linear algebraic group scheme. If F is a field extension of k , the F -rational points of $\mathbf{SL}_1(A)$ are given by $\mathbf{SL}_1(A)(F) = \{x \in A_F \mid \text{Nrd}_{A/F}(a) = 1\}$. Furthermore, by $\mathbf{SK}_1(A)$ we denote the group functor

$$k\text{-fields} \rightarrow \mathbf{Groups} : F \mapsto \mathbf{SK}_1(A)(F) = SK_1(A_F) \cong \mathbf{SL}_1(A)(F)/[A^\times, A^\times],$$

called the *reduced Whitehead group* of A .

Acknowledgements – The author thanks Jean-Louis Colliot-Thélène, Philippe Gille, Bruno Kahn, Marc Levine, and Jean-Pierre Tignol for interesting comments and discussions leading to this article. The author expresses his gratitude to the École Normale Supérieure (Paris), K.U.Leuven, and Research Foundation - Flanders (G.0318.06) for financial support.

2 Existing invariants and results

In this section, we recall the various invariants of $\mathbf{SK}_1$ introduced by several authors. All differ a little bit on the value groups. It takes some time to introduce all of them in quite a rigorous way. The author excuses to the reader who is aware of all the definitions and results and hopes for his tolerance (and endurance). He believes it makes the text more accessible for the non-expert. Experts can eventually skip this section at first and come back to it if necessary to find e.g. a particular definition. Before recalling the invariants, we recall Merkurjev's viewpoint on invariants and Platonov's examples of non-trivial SK_1 as both are used later on.

2.1 Invariants à la Merkurjev

For two group functors $\mathbf{G}, \mathbf{H} : k\text{-fields} \rightarrow \mathbf{Groups}$, an invariant of $\mathbf{G}$ in $\mathbf{H}$ is a natural transformation of functors of $\mathbf{G}$ into $\mathbf{H}$. Typically $\mathbf{H}$ equals the degree j part M_j of a cycle module M (à la Rost [Ros1]), such an invariant is called an *invariant of $\mathbf{G}$ in M of degree j* . It is clear that all invariants of $\mathbf{G}$ in $\mathbf{H}$ form a group (abelian if $\mathbf{H}$ has images in $\mathbf{Ab}$). In case of degree j invariants of $\mathbf{G}$ in a cycle module M , we denote this group by $\text{Inv}^j(\mathbf{G}, M)$. We can define the same terminology if M is any functor of graded groups.

(a) *Cycle modules* – A cycle module M having a field k as base is a formal object having the shared properties of certain Galois cohomology groups, Milnor's K -groups, ... It associates with any field extension F of k a graded abelian group $(M_j)_{j \geq 0}$ endowed with four data (functoriality, reciprocity, K -theory module structure, and residues – D1-D4 in ibid., Def. 1.1) satisfying some homological and geometrical rules (R1a-R3e, FD, and C – ibid., Def. 1.1 & 2.1). For a field k , a central simple k -algebra A of $n = \text{ind}_k(A) \in k^\times$, and an integer $m \in k^\times$, we use the following cycle modules (for any integer r):

$$\mathcal{H}_m^* : k\text{-fields} \mapsto \mathbf{Ab} \quad : \quad F \mapsto (H_m^j(F))_{j \geq 0} \quad (2.1)$$

$$\mathcal{H}_{n,A^{\otimes r}}^* : k\text{-fields} \mapsto \mathbf{Ab} \quad : \quad F \mapsto (H_{n,A^{\otimes r}}^j(F))_{j \geq 1}, \quad (2.2)$$

with

$$H_m^j(F) = H^j(F, \mu_m^{\otimes(j-1)}) \quad \text{and} \quad H_{n,A^{\otimes r}}^j(F) = H_n^j(F) / (H^{j-2}(F, \mu_n^{\otimes(j-2)}) \cup r[A]). \quad (2.3)$$

Remark that if $r \equiv 0 \pmod{\text{per}_k(A)}$, then $H_{n,A^{\otimes r}}^j(F) = H_n^j(F)$. So the second cycle module is actually a generalisation of the first one.

(b) *Gersten complex* – Given a k -variety X and a cycle module M with base k , we have a Gersten complex (ibid., §3.3) (for integers $i, j \geq 0$):

$$\dots \xrightarrow{\partial^{i-2}} \bigoplus_{x \in X^{(i-1)}} M_{j-i+1}(k(x)) \xrightarrow{\partial^{i-1}} \bigoplus_{x \in X^{(i)}} M_{j-i}(k(x)) \xrightarrow{\partial^i} \bigoplus_{x \in X^{(i+1)}} M_{j-i-1}(k(x)) \xrightarrow{\partial^{i+1}} \dots,$$

induced by the residues of the cycle module. Here, $X^{(i)}$ is the set of points of X of codimension i , $k(x)$ is the function field of a point of codimension i , and any appearance of negative degree of the cycle module is to be interpreted as the trivial group. The homology of this complex on spot i is denoted $A^i(X, M_j)$.

(c) *Merkurjev's link* – Let $\mathbf{G}$ be an algebraic k -group which we view as a group functor associating to a field extension F of k , the group $\mathbf{G}(F)$ of F -rational points of $\mathbf{G}$. If M is of bounded exponent, then Merkurjev gives an isomorphism

$$\text{Inv}^j(\mathbf{G}, M) \xrightarrow{\sim} A^0(\mathbf{G}, M_j)_{\text{mult}} \subset A^0(\mathbf{G}, M_j) : \rho \mapsto \rho_K(\xi), \quad (2.4)$$

where $K = k(\mathbf{G})$ is the function field of $\mathbf{G}$ and $\xi \in \mathbf{G}(K)$ the generic point of $\mathbf{G}$ [Mer2, Lem. 2.1 & Thm. 2.3]. The image $A^0(\mathbf{G}, M_j)_{\text{mult}}$ consists of the *multiplicative elements* of $A^0(\mathbf{G}, M_j)$. These are those elements x such that $p_1^*(x) + p_2^*(x) = m^*(x)$ for p_1^*, p_2^*, m^* induced by the projection $p_1, p_2 : \mathbf{G} \times \mathbf{G} \rightarrow \mathbf{G}$ and multiplication $m : \mathbf{G} \times \mathbf{G} \rightarrow \mathbf{G}$.

2.2 Platonov's examples

Among the examples of non-trivial SK_1 of Platonov, we concentrate on the tensor product of two cyclic algebras.

(a) *Cyclic algebras* – Let k be a field and K a cyclic field extension of degree n . Take furthermore a generator $\sigma \in \text{Gal}(K/k) \cong \mathbb{Z}/n$. Then for $b \in k^\times$, we denote by $(K/k, \sigma, b)$ the so-called *cyclic k -algebra* generated by K and a variable x satisfying $x^n = b$ and $xc = \sigma(c)x$ for any $c \in K$. Then clearly $\deg_k(K/k, \sigma, b) = n$ and we can also write this cyclic algebra as $\bigoplus_{i=0}^{n-1} Kx^i$ with multiplication defined as above [Dra, §7, Def. 4]. Furthermore, K is a splitting field of $(K/k, \sigma, b)$ (see [GS, §2.5]).

If k contains an n -th primitive root of unity and if $K = k(\sqrt[n]{a})$ for $a \in k^\times$, then $(K/k, \sigma, b) \cong (a, b)_n$ as k -algebras (if σ is well chosen). Here $(a, b)_n$ is the usual *symbol k -algebra* generated over k by variables x and y satisfying $x^n = b, y^n = a$, and $xy = \xi_n yx$ for a well chosen primitive n -th root of unity $\xi_n \in k$. In case $n = p = \text{char}(k)$ and if K is the cyclic Galois extension defined by $x^p - x = a$, then $(K/k, \sigma, b) \cong [a, b]_p$ as k -algebras (for a well chosen σ). Here $[a, b]_p$ is the usual *p -algebra*: generated as k -algebra by variables x and y satisfying $x^p - x = a, y^p = b$, and $xy = y(x + 1)$ (loc. cit.).

If $n = 2$, a symbol algebra or p -algebra is more commonly called a *quaternion algebra*. The product of two quaternion algebras is a *biquaternion algebra*; it is a central simple algebra of degree 4 and period 1 or 2. It is known that biquaternion algebras are in fact the only central simple algebras of degree 4 and period 1 or 2 [Alb, p. 369].

(b) *Non-trivial SK_1* – Let k be a local field (e.g. $\mathbb{Q}_p$ or $\mathbb{F}_p((x))$) and let K_1, K_2 be two cyclic extensions of degree n over k which are linearly disjoint. Let σ_1 (resp. σ_2) be a generator of $\text{Gal}(K_1/k)$ (resp. $\text{Gal}(K_2/k)$). Now let $F = k((t_1))((t_2))$, $F_1 = K_1((t_1))((t_2))$, and $F_2 = K_2((t_1))((t_2))$. Then Platonov proves that

$$A = (F_1/F, \sigma_1, t_1) \otimes (F_2/F, \sigma_2, t_2)$$

is a division F -algebra and furthermore $SK_1(A) \cong \text{Br}(K/k)/(\text{Br}(K_1/k)\text{Br}(K_2/k)) \cong \mathbb{Z}/n$ for $K = K_1 \otimes K_2$ [Pla, Thms. 4.7 & 5.9].

(c) *Galois cohomology of $\mathbb{Q}_p((t_1))((t_2))$* – To study the invariants later on, we encounter the fourth Galois cohomology groups $H_m^4(k)$ for $k = \mathbb{Q}_p((t_1))((t_2))$. These can be calculated using a splitting for a complete discrete valuation field K with residue field $\kappa(v)$ and with $m \in \kappa(v)^\times$ (hence also $m \in K^\times$) [GMS, 7.11]:

$$H_m^{i+1}(K) \cong H_m^{i+1}(\kappa(v)) \oplus H_m^i(\kappa(v)). \quad (2.5)$$

Using the fact that $\text{cd}(\mathbb{Q}_p) = 2$ and $\text{Br}(\mathbb{Q}_p) = \mathbb{Q}/\mathbb{Z}$ [Ser, Ch. II, §5.1 & Prop. 15], we find $H_m^4(k) \cong \mathbb{Z}/m$ by applying the splitting to the valuations defined by t_1 and t_2 .

2.3 Suslin's invariants

We recall the invariants of Suslin and an invariant for biquaternion algebras introduced by Rost. Let us first give the motivation why these invariants can help to explain Platonov's counterexamples.

(a) *Suslin 1991* – By constructing his invariant $\rho_A \in \text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{m,A}^*)$ (for $m = \text{ind}_k(A) \in k^\times$), Suslin hoped to be able to complete the following diagram (for A as in §2.2 (b)):

$$\begin{array}{ccc} SK_1(A) & \xrightarrow{\cong} & \text{Br}(K/k)/(\text{Br}(K_1/k)\text{Br}(K_2/k)) \\ \rho_{A,F} \downarrow & & \downarrow \\ H_{n^2,A}^4(F) & \xrightarrow{\partial_{t_1}^3 \circ \partial_{t_2}^4} & H_{n^2}^2(k)/\partial_{t_1}^3 \circ \partial_{t_2}^4(H^2(k, \mu_{n^2}^{\otimes 2}) \cup [A]) \end{array} \quad (2.6)$$

The maps $\partial_{t_1}^3, \partial_{t_2}^4$ are residues induced by the discrete valuation associated with t_1 and t_2 , i.e. the projection maps of degree -1 in (2.5). At the time he conjectured the existence of such an invariant, he could not yet give a definition. He was however able to define an invariant $\rho_{\text{S91},A} \in \text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{m,A^{\otimes 2}}^*)$ which he proves to be non-trivial for Platonov's examples of non-trivial SK_1 .

(b) *Biquaternion algebras* – In the case of biquaternion algebras, Rost was able to define a related invariant of $\mathbf{SK}_1(A)$. Suppose A is a biquaternion algebra over a field k of $\text{char}(k) \neq 2$. Then *Rost's invariant* $\rho_{\text{Rost},A}$ is an invariant sitting in $\text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_2^*)$ [Mer1, Thm. 4]. Moreover, it fits into an exact sequence:

$$0 \rightarrow \mathbf{SK}_1(A)(k) \rightarrow H^4(k, \mathbb{Z}/2\mathbb{Z}) \rightarrow H^4(k(Y), \mathbb{Z}/2\mathbb{Z}),$$

where Y an Albert form of A . This invariant was generalised in [KMRT, §17] to biquaternion algebras in any characteristic using Witt groups and Witt rings. We come back to this generalised invariant in Section 3 as its definition requires a lot of terminology related to involutions.

(c) *Suslin 2006* – Using Voevodsky's motivic étale cohomology, Suslin was able to define his conjectured invariant in 2006 [Sus2, §3]. We denote this invariant by $\rho_{\text{S06},A}$. It is however not clear whether (2.6) commutes for this invariant. It is clear that this (and also the other invariants) become trivial after base extension to the function field of $X = \text{SB}(A)$ (it is a splitting field of A). Suslin hence proves his invariant is essentially the same as Rost's invariant $\rho_{\text{Rost},A}$ for a biquaternion algebra over a field k of $\text{char}(k) \neq 2$. He does this by proving that

$$\begin{array}{ccc} \mathbf{SK}_1(A)(k) & \xrightarrow{\rho_{\text{S06}}} & \ker[H_{4,A}^4(k) \rightarrow H_{4,A}^4(k(X))] \\ \downarrow = & & \downarrow r_A \\ \mathbf{SK}_1(A)(k) & \xrightarrow{\rho_{\text{Rost}}} & \ker[H_2^4(k) \rightarrow H_2^4(k(Y))], \end{array} \quad (2.7)$$

is a commutative diagram, where r_A is the morphism induced on Galois cohomology by the map $\mu_4^{\otimes 3} \rightarrow \mu_2 : a \mapsto a^2$ and where X and Y are as above. Hence ρ_{S06} is injective for biquaternion algebras and

$$\mathbf{SK}_1(A)(k) \cong \ker[H_{4,A}^4(k) \rightarrow H_{4,A}^4(k(X))].$$

2.4 Kahn's invariants

Let k be a field and A a central simple algebra with $n = \text{ind}_k(A) \in k^\times$. We recall the inspiring results on invariants of $\mathbf{SK}_1(A)$ as obtained by Kahn in [Kah2].

(a) *Cyclicity of invariant group* – By calculations with motivic étale cohomology, Kahn shows $A^0(\mathbf{SL}_1(A), \mathcal{H}_n^4)_{\text{mult}}$ is finite cyclic [Kah2, Def. 11.3]. So by Merkurjev’s isomorphism (2.4), $\text{Inv}^4(\mathbf{SL}_1(A), \mathcal{H}_n^*)$ is finite cyclic. As the canonical projection $\mathbf{SL}_1(A) \rightarrow \mathbf{SK}_1(A)$ induces an injective morphism

$$\text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_n^*) \rightarrow \text{Inv}^4(\mathbf{SL}_1(A), \mathcal{H}_n^*), \quad (2.8)$$

we also find $\text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_n^*)$ to be cyclic. Using Kahn’s calculations (loc. cit.), we can pick a canonical generator that we call *Kahn’s invariant* $\rho_{\text{Kahn}, A}$ of $\mathbf{SK}_1(A)$.

(b) *Bounds on invariant group* – Kahn also argues the size of $\text{Inv}^4(\mathbf{SL}_1(A), \mathcal{H}_n^*)$ is bounded by $\text{ind}(A)/l$ if $n = \text{ind}_k(A)$ is the power of a prime l (ibid., Lem. 12.1). Hence the same holds for $\text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_n^*)$ by (2.8). For general n , Kahn’s bound is retrieved using Brauer’s decomposition theorem [GS, Ch. 4, Prop. 4.5.16]. For any integer n with prime factorisation $p_1^{e_1} \cdots p_r^{e_r}$, we denote by $\bar{n}$ the integer $p_1^{e_1-1} \cdots p_r^{e_r-1}$.

Lemma 2.1. *Let k be a field and A a central simple algebra of $\text{ind}_k(A) = n \in k^\times$. Then*

$$|\text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_n^*)| \leq \bar{n}.$$

Proof. Let $p_1^{e_1} \cdots p_r^{e_r}$ be a prime decomposition of n , then Brauer’s decomposition theorem (loc.cit.) gives division k -algebras $D_1, \dots, D_r$ of $\text{ind}_k(D_i) = p_i^{e_i}$ such that A is Brauer-equivalent to $D_1 \otimes \dots \otimes D_r$. This even gives rise to a decomposition $\mathbf{SK}_1(A) \cong \mathbf{SK}_1(D_1) \oplus \dots \oplus \mathbf{SK}_1(D_r)$ (ibid., Ch. 4, Ex. 9). Recall also that $\mathbf{SK}_1(D_i)$ has $p_i^{e_i}$ -torsion [Dra, §23, Lem. 3]. Then the result follows immediately from the primary result of Kahn and the isomorphism $H_n^4(k) \cong H_{p_1^{e_1}}^4(k) \oplus \dots \oplus H_{p_r^{e_r}}^4(k)$. ■

Remark 2.2 – As Kahn mentions, this bound is sharp for biquaternion division algebras [Kah2, §12]. This follows from [Mer2, Prop. 4.9 & Thm. 5.4]. In particular, ρ_{Kahn} is not trivial for biquaternion division algebras. In §4.1 (c), we generalise this result.

(c) *Generalisation of Suslin’s invariant* – Apart from using Merkurjev’s viewpoint to define a new invariant, Kahn also generalises ρ_{S06} to invariants

$$\rho_r \in \text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{n, A^{\otimes r}}^*)$$

with $n \in \text{ind}_k(A) \in k^\times$ and $r = 1, \dots, \text{per}_k(A) - 1$. Suslin’s invariant ρ_{S06} is retrieved setting $r = 1$. It is not clear whether ρ_{S91} equals ρ_2 . As mentioned in §2.3 (c), ρ_{S06} has its image in $\ker[H_{n, A}^4(F) \rightarrow H_{n, A}^4(F(X))]$ for F a field extension of k and X the Severi-Brauer variety of A [Sus2, §3]. Kahn generalises this to ρ_r replacing X by the generalised Severi-Brauer variety $\text{SB}(r, A)$ (ibid., §8.B).

He also gives a bound on the torsion of these invariants inside $\text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{n, A}^*)$ if $l = \text{per}_k(A)$ is a prime. Indeed from (ibid., Thm. 7.1(c) & Cor. 12.10) it follows that they have

- l -torsion if $\text{ind}_k(A) = \text{per}_k(A) = l > 2$,
- l^2 -torsion if $\text{per}_k(A) > \text{ind}_k(A) = l > 2$, and
- 2-torsion if $\text{per}_k(A) = 2$.

For a central simple k -algebra A with $n = \text{ind}_k(A) \in k^\times$ and $\text{per}_k(A) = n/\overline{n}$, there is a similar statement using a Brauer decomposition. Take a prime factorisation $n = p_1^{e_1} \cdots p_r^{e_r}$ and let $D_1 \otimes \cdots \otimes D_r$ be a Brauer decomposition of A as in the proof of Lemma 2.1. Then put $m = p_1^{f_1} \cdots p_r^{f_r}$, where $f_i = 1$ if $p_i = 2$ or if $\text{ind}_k(D_i) = \text{per}_k(D_i) = p_i > 2$, and $f_i = 2$ if $\text{ind}_k(D_i) > \text{per}_k(D_i) = p_i > 2$. Then it is clear that ρ_r has m -torsion.

2.5 Generalising invariants

In [Wou], the author introduced a way of generalising the invariants of $\mathbf{SK}_1(A)$ to any central simple k -algebra A (so also when $\text{ind}_k(A) \notin k^\times$). This is done using a lift from a field of positive characteristic to a field of zero characteristic where the invariants are always defined. In this subsection, let k be a field of $\text{char}(k) = p > 0$. We first explain Kato's cohomology of *logarithmic differentials* which are used in (loc. cit.) to generalise $\mathcal{H}_n^*$ when $p \mid n$. This allows us to perform lifts from positive characteristic to characteristic 0.

(a) *Logarithmic differentials* – For any integer $l > 0$, the cohomology groups $H_{p^l}^{q+1}(k)$ are defined as

$$(W_l(k) \otimes \underbrace{k^\times \otimes \cdots \otimes k^\times}_{q \text{ times}}) / I,$$

where $W_l(k)$ are the Witt vectors of length l on k and I is the ideal generated by

- (i) $w \otimes b_1 \otimes \cdots \otimes b_q$, satisfying $b_i = b_j$ for $1 \leq i < j \leq q$,
- (ii) $(0, \dots, 0, a, 0, \dots, 0) \otimes a \otimes b_2 \otimes \cdots \otimes b_q$,
- (iii) $(w^{(p)} - w) \otimes b_1 \otimes \cdots \otimes b_q$,

with $w \in W_l(k)$, $b_1, \dots, b_q \in k^\times$, and $w^{(p)} = (a_1^p, \dots, a_l^p)$ if $w = (a_1, \dots, a_l)$.

For $l = 1$, we can view $H_p^{q+1}(k)$ as the cokernel of

$$F-1 : \Omega_k^q \rightarrow \Omega_k^q / d\Omega_k^{q-1}, \text{ defined by } x \frac{dy_1}{y_1} \wedge \cdots \wedge \frac{dy_q}{y_q} \mapsto (x^p - x) \frac{dy_1}{y_1} \wedge \cdots \wedge \frac{dy_q}{y_q} \pmod{d\Omega_k^{q-1}},$$

hence the terminology “*logarithmic differentials*”. (For $l = 0$, set $H_{p^l}^{q+1}(k) = 0$.)

In general, for an integer $n = p^l m > 0$ ($l, m \geq 0$ integers with $p \nmid m$), we define

$$H_n^{q+1}(k) = H_{p^l}^{q+1}(k) \oplus H_m^{q+1}(k).$$

This is a generalisation of Galois cohomology, since this theory fills in some gaps in Galois cohomology. It gives for example a description of the p^l -th torsion part of the Brauer group, compatible with the prime-to- p part: ${}_p \text{Br}(k) \cong H_{p^l}^2(k)$. So for any integer $n > 0$ we get ${}_n \text{Br}(k) \cong H_n^2(k)$. We can also define $\mathcal{H}_n^*$ in the same way as in (2.1). It is however not a cycle module, but rather a functor of graded groups. To obtain a cycle module we have to tweak it a little bit. For this paper we do not need a cycle module, so we rather work with this functor of graded groups to ease the discussion (see [Wou, §4.1 (d)] for more details – see also Remark 2.6 *infra*).

Using this isomorphism, together with a scalar multiplication by *Milnor's K-groups* on $(H_{p^l}^{q+1}(k))_{q \geq 0}$, we can generalise the definition of $\mathcal{H}_{n,\mathcal{A}}^*$ for a central simple k -algebra

A with arbitrary index. Recall that Milnor K -groups $K_r^M(k)$ (for an integer $r \geq 0$) are defined as

$$\underbrace{k^\times \otimes \dots \otimes k^\times}_{r \text{ times}} / J,$$

where J is the ideal generated by $x_1 \otimes \dots \otimes x_i$ with $x_i + x_j = 1$ for some $1 \leq i < j \leq r$. Elements of $K_r^M(k)$ are called *symbols* and the generators $x_1 \otimes \dots \otimes x_r$ are called *pure symbols*, commonly denoted $\{x_1, \dots, x_r\}$. The scalar multiplication of $K_r^M(k)$ on $(H_{p^l}^{q+1}(k))_{q \geq 0}$ is given by

$$\begin{aligned} \cdot : K_r^M(k) \times H_{p^l}^{q+1}(k) &\rightarrow H_{p^n}^{r+q+1}(k), \quad \text{defined by} \\ (\{x_1, \dots, x_r\}, w \otimes b_1 \otimes \dots \otimes b_q) &\mapsto w \otimes x_1 \otimes \dots \otimes x_r \otimes b_1 \otimes \dots \otimes b_q. \end{aligned}$$

This allows us to define a relative version. Before doing so, we recall that also the cup-product definition of (2.3) can be generalised using K -theory. Indeed, the isomorphism $k^\times / (k^\times)^m \cong H^1(k, \mu_m)$ for any $m \in k^\times$ gives the *Galois symbol* by taking the cup-product:

$$h_{m,k}^r : K_r^M(k) \rightarrow H^r(k, \mu_m^{\otimes r}). \quad (2.9)$$

The Bloch-Kato conjecture (proved by Voevodsky-Rost-Weibel [BK, Voe, Ros2, Wei2]) even says it is surjective with kernel $mK_r^M(k)$. Hence we get a scalar multiplication of $K_r^M(k)$ on $(H_m^{q+1}(k))_{q \geq 0}$:

$$\cdot : K_r^M(k) \times H_m^{q+1}(k) \rightarrow H_m^{r+q+1}(k) \quad \text{defined by} \quad (a, b) \mapsto h_{m,k}^r(a) \cup b.$$

For arbitrary n , this defines in total a $K_r^M(k)$ -module structure on $(H_n^q(k))_{q \geq 0}$. If A is a central simple k -algebra of $\text{ind}_k(A) = n$, we can then define for any field extension F of k and integers $q \geq 0$ and r :

$$H_{n,A^{\otimes r}}^{q+1}(F) = H_n^{q+1}(F) / (K_{q-1}^M(F) \cdot r[A_F]).$$

By the remarks above, this is clearly a generalisation of the moderate case. If $r \equiv 0 \pmod{\text{per}_k(A)}$, then clearly $H_{n,A^{\otimes r}}^{q+1}(F) = H_n^{q+1}(F)$ (cfr. §2.1 (a)). In the same way as in (2.2), we obtain a functor of graded groups $\mathcal{H}_{n,A^{\otimes r}}^*$.

(b) *Lifts* – We now consider k to be the residue field of a complete discrete valuation ring R with fraction field K of $\text{char}(K) = 0$. The specialisation map $\text{Br}(R) \rightarrow \text{Br}(k) : [A] \mapsto [A \otimes_R k]$ is bijective [Gro2, Cor. 6.2] and $\text{Br}(R) \rightarrow \text{Br}(K) : [A] \mapsto [A \otimes_R K]$ is injective [AG, Thm. 7.2]. So we have an inclusion $\text{Br}(k) \rightarrow \text{Br}(K)$; given a central simple algebra A over k , we get a *lifted Azumaya algebra* B over R and an associated central simple algebra B_K over K . Because of the definition, $\text{ind}_k(A) = \text{ind}_K(B_K)$ and by a theorem of Platonov we get $\mathbf{SK}_1(A)(k) \cong \mathbf{SK}_1(B_K)(K)$ [Pla, Thm. 3.12] – see also [Wou, Cor. 3.3].

Furthermore, there exists an injection $H_n^{i+1}(k) \rightarrow H_n^{i+1}(K)$; on the prime-to- p parts of $H_n^{i+1}(k)$ defined by (2.5), for general n see [Kat1, Proof of Prop. 2] and [Izh, Prop. 6.8] (see also Remark 3.7 for 2-primary n). This injection also continues to the relative cohomology groups; i.e. there exists an injection $H_{n,A^{\otimes r}}^{i+1}(k) \rightarrow H_{n,B_K^{\otimes r}}^{i+1}(K)$ for any integer r and A and B as above [Wou, Prop. 4.10].

This allows us to define an invariant for any central simple k -algebra, using the existence in the characteristic 0 case. In order to stay functorial, we have to use p -rings. A

p -ring is a complete discrete valuation ring R with residue field k of $\text{char}(k) = p > 0$ and whose maximal ideal is generated by p . For a reference see e.g. [Mat, §23] where p -rings can also be not complete, we however always suppose them to be complete. For a p -ring R , the fraction field K is of characteristic 0. Moreover, Cohen proved that given a field k of $\text{char}(k) > 0$, there always exists a p -ring with k as residue field [Coh].

For sake of convenience, we also use the following terminology.

Definition 2.3. Suppose ρ is an invariant of $\mathbf{SK}_1$ which is defined for any central simple algebra A with index n not divisible by the characteristic of its base field and which has values in the Galois cohomology group $\mathcal{H}_{n, A^{\otimes r}}^4$ for r a fixed integer. Then we say ρ is a moderate invariant of $\mathbf{SK}_1$ with values in $\mathcal{H}_{\otimes^r}^4$. We denote by ρ_A the invariant for a central simple algebra A .

In [Wou, Thm.4.20], the author proves the following theorem.

Theorem 2.4. Let k be a field of $\text{char}(k) = p > 0$ and A a central simple k -algebra of $n = \text{ind}_k(A)$. Take R a p -ring with residue field k and fraction field K . Let B be the lifted Azumaya R -algebra of A and let $\rho \in \text{Inv}^4(\mathbf{SK}_1(B_K), \mathcal{H}_{n, B_K^{\otimes r}}^*)$ (for r any integer). There exists a unique invariant $\tilde{\rho} \in \text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{n, A^{\otimes r}}^*)$ such that for any field extension k' , p -ring R' with residue field k' , and fraction field K' , we have a commutative diagram:

$$\begin{array}{ccc} \mathbf{SK}_1(A)(k') & \xrightarrow{\tilde{\rho}_{k'}} & H_{n, A^{\otimes r}}^4(k') \\ \cong \uparrow & & \downarrow \\ \mathbf{SK}_1(B_K)(K') & \xrightarrow{\rho_{K'}} & H_{n, B_K^{\otimes r}}^4(K'). \end{array} \quad (2.10)$$

Remark 2.5 – The invariants obtained by this theorem are the *wild generalisations* of their moderate variants (hence the terminology *moderate versus wild*). If ρ is a moderate invariant of $\mathbf{SK}_1$, we denote the wild generalisation by $\tilde{\rho}$. If A is a central simple k -algebra of $\text{ind}_k(A) \in k^\times$ (with $\text{char}(k) = p > 0$), it is in general not clear whether $\rho_A = \tilde{\rho}_A$. By the uniqueness of the theorem, to prove such an equality it suffices to verify that ρ satisfies a lifting property as in (2.10).

Remark 2.6 (for the reader who takes the effort to look at the original paper.) – In the original statement, the author treats just the case $r = 1$. The proof does not depend on r , so it can easily be generalised to any r . If $r = 0$, we can also use (ibid., Cor. 4.14) straightaway to prove the theorem. Also an extra field extension L of k is used. This is to be sure $\mathcal{H}_{n, L}^*$ of (ibid., Def. 4.3) is a cycle module with base R . We do not explicitly need this here. Even more, the statement over here is not weaker as by functoriality any invariant has images in $\mathcal{H}_{n, L, A^{\otimes r}}^*$.

Remark 2.7 – Note that the theorem actually defines an injective morphism

$$\text{Inv}^4(\mathbf{SK}_1(B_K), \mathcal{H}_{n, B_K^{\otimes r}}^*) \rightarrow \text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{n, A^{\otimes r}}^*). \quad (2.11)$$

As the invariants ρ_{Kahn} and ρ_{S06} are non-trivial for biquaternion algebras in characteristics different from 2, this induces their wild generalisations to be non-trivial for biquaternion algebras in characteristic 2.

3 Biquaternion algebras

In [KMRT, §17], Knus-Merkurjev-Rost-Tignol construct an invariant of the reduced Whitehead group of biquaternion algebras in any characteristic. For sake of brevity we call it *KMRT's invariant*. If the characteristic of the base field is not equal to 2, it is known that this invariant essentially equals Suslin's invariant. In this section, we prove in the characteristic 2 case it is essentially equal to Suslin's generalised invariant.

3.1 Definition

We start by giving the concrete definition of KMRT's invariant. This needs the notion of involutions on Azumaya algebras and Witt groups.

(a) *Involutions on Azumaya algebras* – In order to define the invariant, a symplectic involution σ on the biquaternion algebra is used. We recall the definition of a symplectic involution on an Azumaya algebra (so in particular on a central simple algebra). We treat this in this general setting of Azumaya algebras, because we need this for our purposes later on. We refer to [Knu, Ch. III, §8] for more details on involutions on Azumaya algebras.

Definition 3.1. *Let R be a ring and A an Azumaya algebra over R with an R -linear involution σ . Suppose $\alpha : A \otimes_R S \xrightarrow{\sim} M_n(S)$ is a faithfully flat splitting of A . Then $\tilde{\sigma} = \alpha(1 \otimes \sigma)\alpha^{-1}$ is an involution on $M_n(S)$. Since $x \mapsto \tilde{\sigma}(x^t)$ is an automorphism of $M_n(S)$, we can choose $u \in GL_n(S)$ such that $\tilde{\sigma}(x) = ux^tu^{-1}$ for all $x \in M_n(S)$. Because $\tilde{\sigma}^2 = 1$, we get $u^t = \epsilon u$ for some $\epsilon \in \mu_2(S)$. Then ϵ is called the type of σ (it is well defined and independent of the choice of a faithfully flat splitting [Knu, Ch. III, 8.1.1.]). An involution of type 1 is called orthogonal and an involution of type -1 is called symplectic.*

Remark 3.2 – If R is an integral domain, then an involution on an Azumaya algebra can only have type 1 or -1 . When k is a field, a central simple k -algebra of odd degree can only have orthogonal involutions, while a central simple algebra of even degree can have involutions of both types [KMRT, Cor. 2.8].

If A is a central simple algebra over k of degree $2n$ with a symplectic involution σ , we can refine the definition of reduced norm, trace and characteristic polynomial. Indeed, if $a \in \text{Symd}(A, \sigma) = \{a + \sigma(a) \mid a \in A\}$, the reduced characteristic polynomial $\text{Prd}_{A, a/k}(X)$ is a square [KMRT, Prop. 2.9]. Take $\text{Prp}_{\sigma, a/k}(X)$ the unique monic polynomial such that $\text{Prd}_{A, a/k}(X) = (\text{Prp}_{\sigma, a/k}(X))^2$; this is the *Pfaffian characteristic polynomial*. The *Pfaffian trace* $\text{Trp}_{\sigma/k}(a)$ and the *Pfaffian norm* $\text{Nrp}_{\sigma/k}(a)$ are defined as coefficients of $\text{Prp}_{\sigma, a/k}(X)$, compatible with the expression of $\text{Nrd}_{A/k}(a)$ and $\text{Trd}_{A/k}(a)$ as coefficients of $\text{Prd}_{A, a/k}(X)$ (see standard notations in §1):

$$\text{Prp}_{\sigma, a/k}(X) = X^n - \text{Trp}_{\sigma/k}(a)X^{n-1} + \dots + (-1)^n \text{Nrp}_{\sigma/k}(a).$$

So $\text{Nrd}_{A/k}(a) = (\text{Nrp}_{\sigma/k}(a))^2$ and $\text{Trd}_{A/k}(a) = 2\text{Trp}_{\sigma/k}(a)$.

(b) *Witt groups* – To explain the value group of KMRT’s invariant, we need Witt groups and rings.¹ The *Witt group* $W_q(k)$ is the group of Witt-equivalence classes of non-singular quadratic spaces over k with addition defined by the orthogonal sum $\perp$. The *Witt ring* $W(k)$ is the ring of Witt-equivalence classes of non-singular symmetric bilinear spaces with addition given by the orthogonal sum $\perp$ and multiplication by the tensor product $\otimes$.

Remark 3.3 – If $\text{char}(k) \neq 2$, we know that as groups (with the orthogonal sum) $W_q(k)$ and $W(k)$ are isomorphic; not as rings, since one can not come up with a direct definition of multiplication of quadratic forms. For our purposes we are however interested in the characteristic 2 case, so we have to make a clear distinction. For more information on Witt groups and Witt rings in this general case, we refer to [Bae, Ch. I] and [Kah1, Ch. 1] (including the discussion on the characteristic 2 case by Laghribi in [Kah1, App. E]).

We can equip $W_q(k)$ with a $W(k)$ -module structure. If (V, B) is a non-singular symmetric bilinear space on k and (V', q) is a non-singular quadratic space on k , then $(V \otimes V', B \otimes q)$ is a quadratic space on k with $B \otimes q$ defined by

$$(B \otimes q)(v \otimes v') = B(v, v)q(v'), \quad \text{for } v \in V, v' \in V'.$$

Let $I(k)$ be the *fundamental ideal* of $W(k)$ (generated by the non-singular bilinear spaces of even dimension). For any integer $n \geq 0$, we set $I^n(k) = (I(k))^n$ (with $I^0(k) = W(k)$) and $I^n W_q(k) = I^n(k) \otimes W_q(k)$. This clearly defines a filtration

$$W_q(k) = I^0 W_q(k) \supset I^1 W_q(k) \supset I^2 W_q(k) \supset \dots$$

We denote the graded quotients by $\overline{I^n W_q(k)} = I^n W_q(k) / I^{n+1} W_q(k)$.

Remark 3.4 – Set $W'_q(k)$ the subgroup of $W_q(k)$ consisting of equivalence classes of even-dimensional non-singular quadratic spaces over k and $I^n W'_q(k) = I^n(k) \otimes W'_q(k)$. If $\text{char}(k) \neq 2$, we have $I^n W'_q(k) = I^{n+1}(k)$ by the equivalence of symmetric bilinear and quadratic spaces. Again, in general we are not able to use this fact.

(c) *Definition* – Suppose A is a biquaternion algebra over k (see §2.2 (a)) and suppose furthermore that σ is a symplectic involution on A . Knus-Merkurjev-Rost-Tignol construct an explicit map

$$\text{SL}_1(A) \rightarrow \overline{I^3 W'_q(k)} : a \mapsto \begin{cases} 0 & \text{if } \sigma \text{ is hyperbolic,} \\ \Phi_v + I^4 W'_q(k) & \text{if } \sigma \text{ is not hyperbolic,} \end{cases}$$

with kernel equal to $[A^\times, A^\times]$. Recall that an involution is called *hyperbolic* if there exists an idempotent $e \in A$ such that $\sigma(e) = 1 - e$. Furthermore, Φ_v is the quadratic form

$$A \rightarrow k : x \mapsto \Phi_v(x) = \text{Trp}_\sigma(\sigma(x)vx),$$

where $v \in \text{Symd}(A, \sigma) \cap A^\times$ satisfies $v(\text{Trp}_\sigma(v) - v)^{-1} = -\sigma(a)a$. There always exists a v satisfying this condition [KMRT, Lem. 17.3]. This definition is well defined and independent of the choice of v and σ . Moreover the construction is functorial so that we get an invariant $\rho_{\text{BI}, A}$ of $\mathbf{SK}_1(A)$.

¹Do not mix up the Witt group and Witt ring with $W_n(k)$ consisting of the Witt vectors on a field k – see §2.5 (a).

3.2 Comparison, moderate case

In this section, we recall why $\rho_{\text{BI},A}$ and $\rho_{\text{S06},A}$ are equal if A is a biquaternion algebra over k with $\text{char}(k) \neq 2$. This is because both Suslin and Knus-Merkurjev-Rost-Tignol proved their invariant of $\mathbf{SK}_1(A)$ equals $\rho_{\text{Rost},A}$. We already recalled the commutative diagram (2.7) giving us the equality of $\rho_{\text{S06},A}$ and $\rho_{\text{Rost},A}$.

To compare ρ_{BI} to ρ_{Rost} , famous isomorphisms are used, most of them recently proved. Indeed, there are isomorphisms $\psi_F^1 : K_4^M(F)/2 \rightarrow \overline{I^4(F)} = I^4(F)/I^5(F)$ for any F of $\text{char}(F) \neq 2$ (Milnor's conjecture for quadratic forms [Mil, Q. 4.3], proved by Orlov-Vishik-Voevodsky [OVV, Thm 4.1]) and $\psi_F^2 : H^4(F, \mu_2) \rightarrow K_4^M(F)/2$ (Milnor's conjecture [Mil, §6] or a special case of the Bloch-Kato conjecture (2.9)).

So the obvious way of comparing ρ_{BI} and ρ_{Rost} is by the composed isomorphism $\psi_F = \psi_F^1 \circ \psi_F^2$. Indeed, Knus-Merkurjev-Rost-Tignol prove that the following diagram commutes [KMRT, Notes §17]:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbf{SK}_1(A)(F) & \xrightarrow{\rho_{\text{Rost},A,F}} & H_2^4(F) & \longrightarrow & H_2^4(F(Y)) \\ & & \downarrow = & & \downarrow \cong \psi & & \downarrow \cong \\ 0 & \longrightarrow & \mathbf{SK}_1(A)(F) & \xrightarrow{\rho_{\text{BI},A,F}} & \overline{I^4(F)} & \longrightarrow & \overline{I^4(F(Y))}, \end{array} \quad (3.1)$$

for F any field extension of k and Y the Albert form attached to A from §2.3 (b).

So combining (2.7) and (3.1), it follows that ρ_{S06} and ρ_{BI} are the same for biquaternion algebras in characteristic different from 2.

3.3 Comparison, wild case

We first explain how to lift central simple algebras with a symplectic involution. We do this for general central simple algebras and later on use the result for biquaternion algebras.

(a) *Lifting algebras with involution* – Let k be a field of $\text{char}(k) = p > 0$ and R a p -ring with residue field k and fraction field K . Take an Azumaya algebra A over R of degree $2n$ with symplectic involution σ . Define the R -group scheme $\mathbf{PGSp}(A, \sigma) = \mathbf{Aut}(A, \sigma)$, defined for any R -algebra S by

$$\mathbf{Aut}(A, \sigma)(S) = \text{Aut}(A_S, \sigma_S) = \{\varphi \in \text{Aut}_S(A_S) \mid \varphi \circ \sigma_S = \sigma_S \circ \varphi\},$$

with $\sigma_S = \sigma \otimes \text{id}$ the canonical extension of σ to A_S . All Azumaya algebras of degree $2n$ with symplectic involutions up to isomorphism are classified by $H_{\text{ét}}^1(R, \mathbf{PGSp}(A, \sigma))$ [KMRT, 29.22]. Since $\mathbf{PGSp}(A, \sigma)$ is a smooth group scheme (proof as in the field case [KMRT, p. 347]), we can use Hensel's lemma à la Grothendieck to get an isomorphism [SGA, Exp. XXIV, Prop. 8.1]:

$$H_{\text{ét}}^1(R, \mathbf{PGSp}(A, \sigma)) \cong H^1(k, \mathbf{PGSp}(\overline{A}, \overline{\sigma})),$$

where $\overline{A} = A \otimes_R k$ is the reduced central simple k -algebra and $\overline{\sigma} = \sigma \otimes \text{id}$ is the reduced involution on $\overline{A}$, which is also symplectic. On the other hand, we have an inclusion

$$H_{\text{ét}}^1(R, \mathbf{PGSp}(A, \sigma)) \hookrightarrow H^1(K, \mathbf{PGSp}(A_K, \sigma_K)).$$

So in total, we have an inclusion

$$H^1(k, \mathbf{PGSp}(\bar{A}, \bar{\sigma})) \hookrightarrow H^1(K, \mathbf{PGSp}(A_K, \sigma_K)).$$

Remark 3.5 – Note that this lift coincides with lifting central simple algebras as explained in §2.5 (b). Over there we actually used the same arguments for the smooth R -group scheme $\mathbf{PGL}_{R,\infty}$ in order to prove

$$\mathrm{Br}(k) = H^1(k, \mathbf{PGL}_{k,\infty}) \hookrightarrow H^1(K, \mathbf{PGL}_{K,\infty}) = \mathrm{Br}(K).$$

So starting with a central simple k -algebra A with symplectic involution σ , we find a lifted Azumaya algebra B over R with symplectic involution τ and hence a central simple K -algebra B_K with symplectic involution τ_K . In particular, $\deg_k(A) = \deg_K(B_K)$ and $\mathrm{per}_k(A) = \mathrm{per}_K(B_K)$. Since biquaternion algebras are exactly the central simple algebras of degree 4 and period 1 or 2, we see that a biquaternion algebra over k with symplectic involution lifts to a biquaternion algebra with symplectic involution over K .

(b) *Preparing the ingredients* – We now continue the work of §3.2 in the wild case. Throughout this section, let k be a field of characteristic 2, R a 2-ring with residue field k and fraction field K , and A a biquaternion algebra over k with lifted Azumaya algebra B over R . As $\tilde{\rho}_{\mathrm{S06}}$ and ρ_{BI} have different value groups, we first give some remarks on how they relate and how we can use the uniqueness statement of Theorem 2.4 to compare the invariants.

By a theorem of Kato, we have an isomorphism $\psi_k : H_2^4(k) \rightarrow \overline{I^3 W_q(k)}$ [Kat2]. Similar to Suslin's construction (2.7), we can also give a morphism $H_{4,A}^4(k) \rightarrow H_2^4(k)$. Indeed, the *projection*

$$\pi_1^2 : W_2(k) \rightarrow W_1(k) : (a_0, a_1) \rightarrow (a_0)$$

gives a morphism $r : H_4^4(k) \rightarrow H_2^4(k)$. Since π_1^2 sends elements of order 2 to 0, r does exactly the same. Hence we get a morphism $r_A : H_{4,A}^4(k) \rightarrow H_2^4(k)$ because any element of $K_2^M(k) \cdot [A]$ is of order 2. Now we can compare the different groups with a commutative diagram.

Proposition 3.6. *Let k' be a field extension of k and R' a 2-ring (containing R) with residue field k' and fraction field K' , then the following diagram commutes:*

$$\begin{array}{ccccc} H_{4,A}^4(k') & \xrightarrow{r_A} & H_2^4(k') & \xrightarrow[\cong]{\psi_{k'}} & \overline{I^3 W_q(k')} \\ \downarrow i^* & & \downarrow i^* & & \downarrow j \\ H_{4,B_K}^4(K') & \xrightarrow{r_B} & H_2^4(K') & \xrightarrow[\psi_{K'}]{\cong} & \overline{I^3 W_q(K')}. \end{array} \quad (3.2)$$

Remark 3.7 – The morphisms $r_B = r_{B_{K'}}$ and $\psi_{K'}$ are as in (2.7) and (3.1), while $r_A = r_{A_{k'}}$ and $\psi_{k'}$ are as above. The morphism j on Witt groups is as in [Bae, Ch. V, Cor. 1.5]; it is the composition of a bijection of $W_q(R') \cong W_q(k')$ induced by the residual morphism $R' \rightarrow k'$ and an injection $W_q(R') \rightarrow W_q(K')$. Here $W_q(R')$ is the Witt group of quadratic spaces of *constant rank* over R' . See [Bae, Ch. I and V] for more information.

The maps i^* are defined by Kato as in §2.5 (b). We recall the exact definition of this morphism which we need in the proof: for any integer $n > 0$

$$i^* : H_{2^n}^{q+1}(k') \rightarrow H_{2^n}^{q+1}(K') \text{ is defined by } w \otimes \bar{b}_1 \otimes \dots \otimes \bar{b}_q \mapsto i(w) \cup h_{2^n, K'}^q(\{b_1, \dots, b_q\}),$$

where $b_1, \dots, b_q \in R'$, the morphism $h_{2^n, K'}^q$ is the Galois symbol (2.9) and $i(w)$ is the composition

$$W_n(k')/\{w^{(p)} - w | w \in W_n(k')\} \cong H^1(k', \mathbb{Z}/2^n\mathbb{Z}) \xrightarrow{\iota} H^1(K', \mathbb{Z}/2^n\mathbb{Z}),$$

where the isomorphism is induced by the additive form of Hilbert 90 for $W_n(k'_s)$ applied to Witt's short exact sequence [Wit, §5]:

$$0 \longrightarrow \mathbb{Z}/2^n\mathbb{Z} \longrightarrow W_n(k'_s) \xrightarrow{x^{(2)}-x} W_n(k'_s) \longrightarrow 0.$$

The injection ι is defined in a similar way as one can get an injection from the splitting (2.5). It can be proved that i^* behaves well by going to the relative cohomology groups $H_{4,A}^4(k')$ and $H_{4,B_K}^4(K')$ [Wou, Prop. 4.10].

Proof. Let R'_{nr} be a 2-ring with residue field k'_s and fraction field K'_{nr} . So R'_{nr} is the integral closure of R' in K'_{nr} .

We first prove $i^* \circ r_A = r_B \circ i^*$. This follows merely by the definition of i^* . Let $(a_0, a_1) \otimes x_1 \otimes x_2 \otimes x_3 \in H_{4,A}^4(k')$ and take $(b_0, b_1) \in W_2(k'_s)$ such that $(b_0^2, b_1^2) - (b_0, b_1) = (a_0, a_1)$. Then $(a_0) = (b_0)^2 - (b_0) \in W_1(k')$ and

$$i^* \circ r_A((a_0, a_1) \otimes x_1 \otimes x_2 \otimes x_3) = (\bar{\sigma}(b_0) - b_0)_{\sigma \in \Gamma_{K'}} \cup h_2^3(\{x_1, x_2, x_3\}),$$

where we consider $\bar{\sigma}(b_0) - b_0$ as an element of $\mathbb{Z}/2\mathbb{Z}$ for any $\sigma \in \Gamma_{K'}$ (with residue $\bar{\sigma} \in \Gamma_{k'}$). On the other hand,

$$\begin{aligned} r_B \circ i^*((a_0, a_1) \otimes x_1 \otimes x_2 \otimes x_3) &= r_B[(\bar{\sigma}(b_0, b_1) - (b_0, b_1))_{\sigma \in \Gamma_{K'}} \cup h_4^3(\{x_1, x_2, x_3\})] \\ &= (\bar{\sigma}(b_0) - (b_0))_{\sigma \in \Gamma_{K'}} \cup h_2^3(\{x_1, x_2, x_3\}). \end{aligned}$$

The commutativity of the right square is essentially due to Kato [Kat2, Lem. 11]; he proves the existence of a commutative diagram

$$\begin{array}{ccc} H_2^n(k') & \xrightarrow{\cong} & \overline{I^3 W_q(k')} \\ \downarrow \varphi & & \downarrow j \\ K_n^M(K')/2K_n^M(K') & \xrightarrow[\psi_{K'}^1]{\cong} & \overline{I^3 W_q(K')} \end{array}$$

where $\psi_{K'}^1$ is the isomorphism of Milnor's conjecture on quadratic forms (see §3.2) and φ is defined by

$$\bar{b} \frac{d\bar{a}_1}{\bar{a}_1} \wedge \frac{d\bar{a}_2}{\bar{a}_2} \wedge \frac{d\bar{a}_3}{\bar{a}_3} \mod I \mapsto \{1 + 4b, a_1, a_2, a_3\} \mod 2K_n^M(K'),$$

for $a_1, a_2, a_3, b \in R'$. Since the isomorphism $\psi_{K'}^1 : H_2^4(K') \rightarrow \overline{I^3 W_q(K')}$ is defined as composition of $\psi_{K'}^1$ with the Galois symbol $h_{2, K'}^4$, it suffices to check $i(\bar{b}) = h_{2, k'}^1(4b + 1)$ for any $b \in R'$. So take $c \in k'_s$ such that $c^2 - c = \bar{b}$. Then

$$i(\bar{b}) = (\bar{\sigma}(c) - c)_{\sigma \in \Gamma_{K'}} \in H^1(K', \mathbb{Z}/2)$$

under the standard identification of $\mathbb{Z}/2$ and $\mu_2(K')$. Take $\tilde{c}$ to be a lift of c in R_{nr} . By eventually changing the representant of $\bar{b}$ in R' , we can assume $\tilde{c}^2 - \tilde{c} = b$. Then $4b + 1 = (2\tilde{c} + 1)^2$ and

$$h_{2, K'}^1(4b + 1) = (\sigma(2\tilde{c} + 1)/(2\tilde{c} + 1))_{\sigma \in \Gamma_{K'}} \in H_2^1(K').$$

So if $\sigma(2\tilde{c} + 1)/(2\tilde{c} + 1) = 1$, we have $\sigma(\tilde{c}) = \tilde{c}$. On the other hand, if $\sigma(2\tilde{c} + 1)/(2\tilde{c} + 1) = -1$, we get $\sigma(\tilde{c}) = -\tilde{c} - 1$. This gives indeed the desired equality. $\blacksquare$

(c) *Cooking up the result* – Using Theorem 2.4 and Proposition 3.6, we can prove the main theorem.

Theorem 3.8. *Let k be a field of characteristic 2 and A a biquaternion algebra over k , then for any field extension k' of k*

$$\rho_{\text{BI}, A, k'} = \psi_{k'} \circ r_A \circ \tilde{\rho}_{\text{S06}, A, k'}$$

with $\psi_{k'}$ and r_A as in (3.2).

Proof. Let k' be a field extension of k and R (resp. R') a 2-ring with residue field k (resp. k') and fraction field K (resp. K'). Suppose σ is a symplectic involution on A and take B a lifted Azumaya R -algebra with lifted symplectic involution τ . Use the same notations as in (3.2). We know j is injective, $i^* \circ \tilde{\rho}_{\text{S06}, A} = \rho_{\text{S06}, B_K}$ (by definition of $\tilde{\rho}_{\text{S06}, A}$) and $\rho_{\text{BI}, B_K} = \psi_{K'} \circ r_B \circ \rho_{\text{S06}, B_K}$. So it suffices to prove that $\rho_{\text{BI}, B_K} = j \circ \rho_{\text{BI}, A}$, which merely follows from the definition.

Let us first explain the isomorphism $\mathbf{SK}_1(B_K)(K') \cong \mathbf{SK}_1(A)(k')$. We can suppose that $\mathbf{SK}_1(A)(k') \neq 0$ so that $A_{k'}$ and $B_{K'}$ are division algebras by Wang's theorem [Wan]. Then $B_{K'}$ is equipped with a valuation w that extends the valuation v' of K' , namely $\frac{1}{4}v' \circ \text{Nrd}_{B_{K'}/K'}$. The associated valuation ring is $B_{R'}$ and the reduced k -algebra is $A_{k'}$. Even more, $\text{SL}_1(B_{K'})$ is part of $B_{R'}$ and the isomorphism $\mathbf{SK}_1(B_K)(K') \cong \mathbf{SK}_1(A)(k')$ is induced by the residue map on $\mathbf{SL}_1(B_{K'})$ [Pla, Cor. 3.13] – see also [Wou, Cor. 3.3].

The involutions σ and τ can not be hyperbolic due to [KMRT, Prop. 6.7 (3)]. Take $a \in \mathbf{SK}_1(A)(k')$ with lift $b \in \mathbf{SK}_1(B_K)(K')$. Then by definition it follows that $\text{Prd}_{A_{k'}, a/k'}(X) = \overline{\text{Prd}_{B_{K'}, b/K'}(X)}$, where the residue is the canonical residue on $R'[X]$. So we also get $\text{Prp}_{\sigma_{k'}, a/k'}(X) = \overline{\text{Prp}_{\tau_{K'}, b/K'}(X)}$ and $\text{Trp}_{\sigma_{k'}/k'}(a) = \overline{\text{Trp}_{\tau_{K'}/K'}(b)}$. Then take $y \in \text{Symd}(B_{K'}, \tau_{K'}) \cap B_{K'}^\times$, satisfying $y(\text{Trp}_{\tau_{K'}/K'}(y) - y)^{-1} = -\tau(b)b$. We can assume $w(y) \geq 0$, since if $w(y) < 0$, i.e. $\text{Nrd}_{B_{K'}/K'}(y) = \lambda/\mu \in K'$ with $\lambda, \mu \in R'$, then $w(\mu y) = v(\lambda) \geq 0$ and

$$\mu y \left(\text{Trp}_{\tau_{K'}/K'}(\mu y) - \mu y \right)^{-1} = y(\text{Trp}_{\tau_{K'}/K'}(y) - y)^{-1}.$$

Then we get $\bar{y}(\text{Trp}_{\sigma_{k'}/k'}(\bar{y}) - \bar{y})^{-1} = -\sigma(a)a$ as b is a lift of a and moreover $\bar{y} \in \text{Symd}(A, \sigma)$. Hence

$$\begin{aligned} \rho_{\text{BI}, A, k'}(a) &= \Phi_{\bar{y}} : A_{k'} \rightarrow k' : x \mapsto \text{Trp}_{\sigma_{k'}/k'}(\sigma_{k'}(x)\bar{y}x) \quad \text{and} \\ \rho_{\text{BI}, B_{K'}, K'}(b) &= \Phi_y : B_{K'} \rightarrow K' : x \mapsto \text{Trp}_{\tau_{K'}/K'}(\tau_{K'}(x)y x). \end{aligned}$$

As $\overline{\text{Trp}_{\tau_{K'}/K'}(\tau_{K'}(x)y x)} = \text{Trp}_{\sigma_{k'}/k'}(\sigma_{k'}(\bar{x})\bar{y}\bar{x})$ for $x \in B_{R'}$, the required compatibility holds. $\blacksquare$

3.4 Non-triviality of the invariant

Because the invariants for biquaternions in odd or zero characteristic are injective, they are also injective in characteristic 2 due to the lifting property (Theorem 2.4). As $\mathbf{SK}_1$ is not trivial for Platonov's examples (§2.2 (b)) and in general for biquaternion algebras of index 4 [Mer3], we find non-trivial invariants in characteristic 2.

Another argument for non-triviality of ρ_{BI} in characteristic different from 2 is given by a formula of Merkurjev for the value on the centre of the biquaternion algebra [Mer1, Ex. p. 70] – see also [KMRT, Ex. 17.23]. Using this formula and the lift from characteristic 2 to characteristic 0, one could hope to prove the non-triviality of ρ_{BI} (and hence of ρ_{S06}) in the case when $\text{char}(k) = 2$, but this fails. Let us comment on this fact.

Say k is a field of characteristic 2, R a p -ring with residue field k and fraction field K , and let $A = [\bar{a}, \bar{b}] \otimes_k [\bar{c}, \bar{d}]$ be a biquaternion k -algebra for $a, c \in R$ and $b, d \in R^\times$. Then the lifted Azumaya R -algebra is $B = [a, b] \otimes_R [c, d]$ where e.g. $[a, b]$ is the R -algebra generated by u, v satisfying slightly different relations than usual: $u^2 + u = a$, $v^2 = b$, and $uv = -v(u + 1)$. We can rewrite it as $B = (4a + 1, b)_R \otimes_R (4c + 1, d)_R$, where $(4a + 1, b)_R$ is the R -algebra generated by i, j with $i^2 = 4a + 1$, $j^2 = b$, and $ij = -ji$. Indeed, an isomorphism is given by $i = 2u + 1$ and $j = v$. Suppose K contains a primitive fourth root of unity ζ , then by (loc. cit.) we have

$$\rho_{\text{BI}, B_K, K}([\zeta]) = \langle\langle 4a + 1, b, 4c + 1, d \rangle\rangle + I^4 W'_q(K),$$

where $[\zeta]$ is the class of ζ in $\mathbf{SK}_1(B_K)(K)$ and where $\langle\langle 4a + 1, b, 4c + 1, d \rangle\rangle$ is an n -fold Pfister quadratic K -form [Kah1, Lem. 2.1.1].

Let π be the isomorphism $\mathbf{SK}_1(B_K)(K) \cong \mathbf{SK}_1(A)(k)$, then $\pi([\zeta]) = [1]$ because k contains no non-trivial fourth roots of unity. By the proof of Theorem 3.8, we have $j \circ \rho_{\text{BI}, B_K, K}([\zeta]) = \rho_{\text{BI}, A, k} \circ \pi([\zeta]) = 0 \in \overline{I^3 W'_q(k)}$. Because the map j from Proposition 3.6 is injective, we get that $\langle\langle 4a + 1, b, 4c + 1, d \rangle\rangle = 0 \in \overline{I^3 W'_q(K)}$. We can also verify this by calculating with Pfister forms. Define $\mathcal{Q} = (4a + 1, b)_R$ and let $\mathcal{X}$ be the natural affine R -scheme with

$$\mathcal{X}(R) = \{x \in \mathcal{Q} \mid \text{Nrd}_{\mathcal{Q}_K/K}(x) = 4c + 1\},$$

where $\mathcal{Q}_K = \mathcal{Q} \otimes_R K$. Then $\mathcal{X}$ is an R -torsor under $\mathbf{SL}_1(\mathcal{Q})$, where $\mathbf{SL}_1(\mathcal{Q})$ is the natural affine R -scheme so that $\mathbf{SL}_1(\mathcal{Q})(R) = \mathbf{SL}_1(\mathcal{Q}_K)(K) \cap \mathcal{Q}$. The special fibre $\mathcal{X}_k = \mathcal{X} \times_R k$ clearly has a rational point, so its class $[\mathcal{X}_k] \in H^1(k, \mathbf{SL}_1(\mathcal{Q}_k))$ is trivial. By Hensel's lemma [SGA, Exp. XXIV, Prop. 8.1], we get $[\mathcal{X}] = 0 \in H_{\text{ét}}^1(R, \mathbf{SL}_1(\mathcal{Q}))$. Hence $\mathcal{X}$ (as well as the generic fibre $\mathcal{X}_K$) has a rational point, but then by theory of Pfister forms we get $\langle\langle 4a + 1, b, 4c + 1 \rangle\rangle = 0 \in W'_q(K)$ [Kah1, Cor. 2.1.10]. Indeed, $\text{Nrd}_{\mathcal{Q}_K/K}(x)$ corresponds with a value of $\langle\langle 4a + 1, b \rangle\rangle$. So a fortiori $\langle\langle 4a + 1, b, 4c + 1, d \rangle\rangle = 0 \in \overline{I^3 W'_q(K)}$.

4 Comparing to Kahn's invariant

We compare now all defined invariants of $\mathbf{SK}_1(A)$ to $\rho_{\text{Kahn}, A}$ in the moderate case, i.e. as they are originally defined. The results can be generalised to the wild invariants, but with some loss of information. We also generalise the formula of Merkurjev for the value on the centre of $\mathbf{SK}_1(A)$ (§3.4).

4.1 Moderate case

We explain two natural ways of comparing $\text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_n^*)$ and $\text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{n, A^{\otimes r}}^*)$. Let A be a central simple k -algebra with $\text{ind}_k(A) = n \in k^\times$ and $m = \text{per}_k(A)$.

(a) *Ways of looking* – For any field extension F of k and any integer r , we can look at the composition

$$m_r : H_{n, A^{\otimes r}}^4(F) \xrightarrow{m} H_{n/m}^4(F) \hookrightarrow H_n^4(F)$$

and at the projection

$$\pi_r : H_n^4(F) \rightarrow H_{n, A^{\otimes r}}^4(F).$$

These induce respectively maps

$$\begin{aligned} \tilde{m}_r : \text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{n, A^{\otimes r}}^*) &\rightarrow \text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_n^*) \quad \text{and} \\ \tilde{\pi}_r : \text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_n^*) &\rightarrow \text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{n, A^{\otimes r}}^*). \end{aligned}$$

The maps $\tilde{\pi}_r$ were introduced by Kahn [Kah2, Rem. 11.6], but we rather consider the maps $\tilde{m}_r$ to compare because of the special definition of Kahn's invariant as generator of the target group. We could also refine $\tilde{m}_r$ if $H^2(k, \mu_n^{\otimes 2}) \cup r[A]$ has m' -torsion for an integer $0 \leq m' < m$. A good comprehension of both maps actually relies, as Kahn mentions, on a good comprehension of the cup product with the class of A (loc. cit.).

By the cyclicity of $\text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_n^*)$ (§2.4 (a)), we certainly find the following relations.

Proposition 4.1. *Let A be a central simple k -algebra of $\text{ind}_k(A) \in k^\times$. Then for any integer r and any $\rho \in \text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{n, A^{\otimes r}}^*)$ there exists an integer $d_A \in \mathbb{Z}/\overline{n}$ such that*

$$\tilde{m}_r(\rho) = d_A \rho_{\text{Kahn}, A} \in \text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_n^*) \subset \mathbb{Z}/\overline{n}.$$

Proof. Use the definition of ρ_{Kahn} and the bounds on $\text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_n^*)$ (see §2.4 (b)). ■

Kahn also raises the issue whether $\tilde{\pi}_r$ is surjective or not (loc. cit.). We can prove it to be non-surjective for biquaternion division algebras à la Platonov.

Proposition 4.2. *Let $k = \mathbb{Q}_p((t_1))((t_2))$ for a prime p . Suppose $A = (a, t_1) \otimes (b, t_2)$ is a biquaternion division k -algebra for $a, b \in \mathbb{Q}_p^\times$. Then $\tilde{\pi}_1$ is not surjective.*

Proof. In §§2.2 (b) and (c) we saw that $SK_1(A) \cong \mathbb{Z}/2$ and $H_4^4(k) \cong \mathbb{Z}/4$. We can also add a fourth primitive root of unity to k as this does not change the Brauer group. In this case we have the Bloch-Kato isomorphism $H_4^4(k) \cong K_4^M(k)/4$.

We now prove $H_{4, A}^4(k) \cong \mathbb{Z}/2$. Under the Bloch-Kato-isomorphism $K_2^M(k)/2 \cong {}_2\text{Br}(k)$, the class of A corresponds to $\{a, t_1\} + \{b, t_2\} \in K_2^M(k)/2$ [GS, Prop. 4.7.1] so that $H^2(k, \mu_4^{\otimes 2}) \cup [A]$ is isomorphic to $(K_2^M(k)/4) \cdot (2\{a, t_1\} + 2\{b, t_2\})$. As the isomorphism $H_4^4(k) \cong \mathbb{Z}/4$ is retrieved by taking two residues $\partial_{t_2}^3$ and $\partial_{t_1}^4$, it suffices to determine the group

$$\partial_{t_2}^3 \circ \partial_{t_1}^4 ((K_2^M(k)/4) \cdot (2\{a, t_1\} + 2\{b, t_2\})).$$

By the definition of residues on Milnor K -groups [Mil, §2], it is clear that this equals $(K_1^M(k)/4) \cdot 2\{a\} + (K_1^M(k)/4) \cdot 2\{b\}$. As we assumed that $\mathbf{SK}_1(A)$ is not trivial, a can not be a square otherwise A would have been Brauer-trivial. This means that $(K_1^M(k)/4) \cdot 2\{a\} + (K_1^M(k)/4) \cdot 2\{b\}$ is not trivial. On the other hand it has 2-torsion inside $K_2^M(k)/4 \cong \mathbb{Z}/4$ so that indeed $H_{4, A}^4(k) \cong \mathbb{Z}/2$.

Then $\pi_1 : \mathbb{Z}/4 \rightarrow \mathbb{Z}/2$ is the “modulo 2” map and $m_1 : \mathbb{Z}/2 \rightarrow \mathbb{Z}/4$ is canonical injection. Suslin proves $\rho_{\text{S06}, A, k} : \mathbf{SK}_1(A)(k) \rightarrow H_{4, A}^4(k)$ is not trivial (2.7), so it is the identity map on $\mathbb{Z}/2$. It is then clear that this can never factor through $H_4^4(k)$ so that $\tilde{\pi}_1$ is clearly not surjective. ■

(b) *Determining factors* – We prove that for the product of two symbol algebras of degree n the factor d_A appearing in Proposition 4.1 only depends on the invariant ρ and the characteristic of k .

Proposition 4.3. *Let ρ be a moderate invariant of $\mathbf{SK}_1$ with values in $\mathcal{H}_{\otimes r}^4$. Let furthermore p be equal to zero or to any prime and let m be an integer not divisible by p . Then there exist an integer $i(p, m) \in \mathbb{Z}/\overline{m^2}$ such that for any field k of $\text{char}(k) = p$ containing a primitive m -th root of unity ξ_m and for any product $A = (a, b)_m \otimes (c, d)_m$ of two symbol k -algebras*

$$\tilde{m}_r(\rho_A) = i(p, m) \rho_{\text{Kahn}, A} \in \text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{m^2}^*) \subset \mathbb{Z}/\overline{m^2}.$$

Remark 4.4 – Although $i(p, m)$ is in general not uniquely determined, we can take a canonical representant as we know $\text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{m^2}^*)$ is cyclic. This comes down to taking the class in $\mathbb{Z}/\overline{m^2}$ satisfying the required relation and such that the representant in $\{0, \dots, m^2 - 1\}$ is as low as possible. It also of course depends on the invariant. We add an index if necessary to stress which invariant is compared to Kahn's invariant. Moreover, it also depends on the exact definition of the injection $\text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{m^2}^*) \subset \mathbb{Z}/\overline{m^2}$. For the remainder of the paper, we fix this injection.

Proof. Take k the prime field of characteristic p and set $k' = k(\xi_m)$ for an m -primitive root of unity $\xi_m \in k_s$. Denote by $\mathcal{T} = (t_1, t_2)_m \otimes (t_3, t_4)_m$ the product of two Azumaya symbol algebra over $R = k'[t_1^{\pm 1}, t_2^{\pm 1}, t_3^{\pm 1}, t_4^{\pm 1}]$ where t_1, t_2, t_3, t_4 are variables and where Azumaya symbol algebras are defined using the same relations as used for symbol algebras over a field. Take $K = k'(t_1, t_2, t_3, t_4)$ and $T = \mathcal{T}_K = (t_1, t_2)_m \otimes (t_3, t_4)_m$, the product of the respective symbol algebras over K . By Proposition 4.1, we find a unique $d_T \in \mathbb{Z}/\overline{m^2}$ such that

$$\tilde{m}_r(\rho_T) = d_T \rho_{\text{Kahn}, T}. \quad (4.1)$$

We prove d_T only depends on m and p .

So suppose F is a field of characteristic p containing an m -th primitive root of unity so that $k' \subset F$. Take any product $A = (a, b)_m \otimes (c, d)_m$ of two symbol algebras of degree m over F . Now A can be obtained from $\mathcal{T}_F = \mathcal{T} \otimes_R F$ by specialising t_1, t_2, t_3, t_4 to a, b, c, d respectively.

Furthermore, (a, b, c, d) defines a k -rational point x of $\text{Spec}(F[t_1^{\pm 1}, t_2^{\pm 1}, t_3^{\pm 1}, t_4^{\pm 1}])$. Take $\mathcal{O}_x$ to be the local ring of $\text{Spec}(F[t_1^{\pm 1}, t_2^{\pm 1}, t_3^{\pm 1}, t_4^{\pm 1}])$ in x with maximal ideal M . It is clear that the completion $\hat{\mathcal{O}}_x$ of $\mathcal{O}_x$ with respect to the M -adic topology is F -isomorphic to $R' = F[[u_1, u_2, u_3, u_4]]$ where $u_1 = t_1 - a, u_2 = t_2 - b, u_3 = t_3 - c$, and $u_4 = t_4 - d$ (see also [Gro1, Thm. 19.6.4]). Under the isomorphism $\text{Br}(R') \cong \text{Br}(F)$ from §2.5 (b), it is clear that $A_{R'} = A \otimes R'$ is an Azumaya R' -algebra mapping to A . Furthermore, the F -isomorphism of $\hat{\mathcal{O}}_x$ with R' gives an isomorphism $\text{Br}(\hat{\mathcal{O}}_x) \cong \text{Br}(R')$. In its turn, this gives an isomorphism $\text{Br}(\hat{\mathcal{O}}_x) \rightarrow \text{Br}(F)$ with inverse given by taking the tensor product over F with $\hat{\mathcal{O}}_x$. By construction it sends the class of $\mathcal{T}_{\hat{\mathcal{O}}_x}$ to the class of A .

Let $K' = F((u_1))((u_2))((u_3))((u_4))$, then $A \otimes_F K'$ is Brauer-equivalent to $\mathcal{T}_{\hat{\mathcal{O}}_x} \otimes_{\hat{\mathcal{O}}_x} K' \cong T_{K'}$. We find $SK_1(A_{K'}) \cong SK_1(T_{K'})$ (as in §2.5 (b)). Furthermore, (2.5) gives an injection

$H_{m^2}^4(F) \rightarrow H_{m^2}^4(K')$. By functoriality of invariants, the diagram

$$\begin{array}{ccc} SK_1(A) & \xrightarrow{\rho} & H_{m^2}^4(F) \\ \cong \downarrow & & \downarrow \\ SK_1(T_{K'}) & \xrightarrow{\rho} & H_{m^2}^4(K') \end{array}$$

commutes both for $\tilde{m}_r(\rho)$ and ρ_{Kahn} . Then by (4.1), we get $\tilde{m}_r(\rho_A) = d_T \rho_{\text{Kahn}, A}$. $\blacksquare$

In particular, we find such relations for $\rho = \rho_{\text{S91}}, \rho_{\text{S06}}$, and the ρ_r 's.

(c) *Non-triviality of Kahn's invariants* – As mentioned in Remark 2.2, ρ_{Kahn} is not-trivial for biquaternion algebras (of index 4). We generalise this to the product of two cyclic algebras à la Platonov (§2.2). Therefore, we compare ρ_{Kahn} to ρ_{S91} as this invariant is non-trivial for Platonov's examples (§2.3 (a)). This means we have to work with $\mathcal{H}_{n, A^{\otimes 2}}^*$ for suitable n and A . (In the same way as in Proposition 4.2, these give also examples of non-trivial $\tilde{\pi}_2$.)

Theorem 4.5. *Let k be p -adic field containing a n^3 -th primitive root unity. Suppose $A = (a, t_1)_n \otimes (c, t_2)_n$ is a division $k((t_1))((t_2))$ -algebra, then $\rho_{\text{Kahn}, A}$ is not trivial. If $n = q_1 \cdot \dots \cdot q_r$ for different primes q_i , then*

$$\text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{n^2}^*) \cong \mathbb{Z}/n.$$

Moreover if n is odd, the integer $i_{\text{S91}}(0, n) \in \mathbb{Z}/\overline{n^2}$ defined in Proposition 4.3 for ρ_{S91} is not trivial.

Proof. We know $SK_1(A) \cong \mathbb{Z}/n$ by §2.2. Furthermore $H_{n^2}^4(k) = \mathbb{Z}/n^2$ as the results in §2.2 (c) hold also when one replaces $\mathbb{Q}_p$ by a finite extension of it.

To calculate $H_{n^2, A^{\otimes 2}}^4(k)$, we use a analogous argument as in the proof of Proposition 4.2. If n is odd, we also find $H_{n^2, A^{\otimes 2}}^4(k) = \mathbb{Z}/n$ as in this case $\text{per}_k(A^{\otimes 2}) = \text{per}_k(A)$. If n is even, $\text{per}_k(A^{\otimes 2}) = n/2$ so that $H_{n^2, A^{\otimes 2}}^4(k) = \mathbb{Z}/(2n)$. In either case, $m_2 : H_{n^2, A^{\otimes 2}}^4(k) \rightarrow H_{n^2}^4(k)$ is the canonical injection (m_2 is the multiplication by m for $m = n$ if n odd and $m = n/2$ if n even).

Suslin proves $\rho_{\text{S91}, A}$ is not trivial (on the field k) [Pla, Thm. 4.8]. If n is odd, $\rho_{\text{Kahn}, A}$ is not trivial (on k) by Proposition 4.1 and hence by definition $i_{\text{S91}}(0, n^2) \neq 0 \in \mathbb{Z}/\overline{n^2}$. If n is even, a similar argument as in the proof of Proposition 4.1 gives the non-triviality of $\rho_{\text{Kahn}, A}$ (mutatis mutandis m by $n/2$).

By the bound on the invariant group (§2.4 (b)) and a Brauer decomposition of A with a related decomposition of invariants in primary parts, the isomorphism statement follows. $\blacksquare$

4.2 Wild case

We continue the comparison in the wild case. Using a lift, we can generalise the statement to any central simple algebra with some loss of information. This does let us prove a relation between the several $i(p, n)$'s.

Let A be a central simple k -algebra of $\text{ind}_k(A) = n$ and $\text{per}_k(A) = m$. We again have morphisms for any integer r

$$\tilde{m}_r : \text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{n, A^{\otimes r}}^*) \rightarrow \text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_n^*)$$

induced by the multiplication for any field extension F of k :

$$m_r : H_{n, A^{\otimes r}}^4(F) \xrightarrow{\cdot m} H_{n/m}^4(F) \hookrightarrow H_n^4(F).$$

Note that we can also define maps $\tilde{\pi}_r$ as in §4.1 (a).

Proposition 4.6. *Let ρ be a moderate invariant of $\mathbf{SK}_1$ with values in $\mathcal{H}_{\otimes r}^4$. Suppose k is a field of $\text{char}(k) = p > 0$ and let $A = [a, b]_p \otimes [c, d]_p$ be the product of two p -algebras over k , then*

$$\tilde{m}_r(\tilde{\rho}_A) = i(0, p)\tilde{\rho}_{\text{Kahn}, A}.$$

Proof. Take a p -ring R with residue field k and fraction field K . Remark first that the lifted Azumaya R -algebra B of A is (after base extension to K) a product of two symbol algebras of degree p . This follows from [GS, Prop. 4.7.1 & Prop. 9.2.5] and the injection $H_{p^2}^2(k) \rightarrow H_{p^2}^2(K)$ [Kat1, Proof of Thm. 1 & 3].

The result follows immediately from the injections

$$\begin{aligned} \text{Inv}^4(\mathbf{SK}_1(B_K), \mathcal{H}_{p^2}^*) &\rightarrow \text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{p^2}^*) \quad \text{and} \\ \text{Inv}^4(\mathbf{SK}_1(B_K), \mathcal{H}_{p^2, B_K^{\otimes r}}^*) &\rightarrow \text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_{p^2, A^{\otimes r}}^*) \end{aligned}$$

defined by lifting invariants and the relations for ρ_{B_K} and ρ_{Kahn, B_K} (Proposition 4.3). $\blacksquare$

Remark 4.7 – In the view of Remark 2.5, we could even refine the statement in the moderate case. If k is a field of $\text{char}(k) = p > 0$ and if $A = (a, b)_n \otimes (c, d)_n$ is the product of two symbol algebras for $n \in k^\times$, then a similar statement holds as A lifts to the product of two symbol algebras of degree n in characteristic 0. If $\tilde{\rho}_A = \rho_A$, then $i(p, n)$ is a multiple of $i(0, n)$ in $\mathbb{Z}/\overline{n}$. Indeed, $\rho_{\text{Kahn}, A}$ is a generator of $\text{Inv}^4(\mathbf{SK}_1(A), \mathcal{H}_n^*) \subset \mathbb{Z}/\overline{n}$ and for some integer d

$$i(p, n)\rho_{\text{Kahn}, A} = \tilde{m}_r(\rho_A) = i(0, n)\tilde{\rho}_{\text{Kahn}, A} = i(0, n)d\rho_{\text{Kahn}, A}.$$

4.3 Formula on the centre

We can now generalise the formula of Merkurjev on the centre of a biquaternion algebra ([Mer1, Ex. p.70] – see also [KMRT, Ex. 17.23] and §3.4) to the tensor product of two symbol algebras. We first prove a general formula and later we prove a finer result using Theorem 4.5.

(a) *General result* – We again use cohomological invariants, however not invariants of algebraic groups as in §2.1, but rather invariants as introduced in [GMS, Ch. I]. These are also natural transformations of functors, but rather a transformation of a functor $B : k\text{-fields} \rightarrow \mathbf{Sets}$ into a functor $H : k\text{-fields} \rightarrow \mathbf{Groups}$.

Proposition 4.8. *Let p be equal to 0 or to any prime and let $n > 0$ be an integer not divisible by p . Let φ be the canonical map $H_m^4(k) \rightarrow H_{n^2}^4(k)$ (for $m = \overline{n^2}$). There exists an integer $j(p, n)$ such that the following formula holds for any field k of $\text{char}(k) = p$ containing a primitive n^2 -th root of unity ζ and for any product $A = (a, b)_n \otimes (c, d)_n$ of two symbol k -algebras:*

$$\rho_{\text{Kahn}, A, k}([\zeta]) = \varphi [j(p, n) h_m^4(\{a, b, c, d\})] \in H_{n^2}^4(k)$$

Remark 4.9 – Remark that as k contains an n^2 -th primitive root of unity, $\mu_{n^2}^{\otimes i} \cong \mathbb{Z}/n^2$ for any $i > 0$. Note also that $\varphi[h_m^4(\{a, b, c, d\})] = (n/m) h_{n^2}^4(\{a, b, c, d\})$.

This expression is compatible to the biquaternion case keeping in mind diagrams (2.7) and (3.1). Also, the integer $j(p, n)$ in the theorem is not uniquely determined, but can be picked canonically by taking the smallest positive integer satisfying the relation. Moreover, $j(p, n)$ depends on the n -th primitive root of unity used in the definition of the symbol algebra and of the choice of n^2 -th primitive root of unity ζ . We are interested in the invertibility of $j(p, n)$ modulo m and therefore the exact choices do not matter, so we do not incorporate these in the notation.

Proof. As ρ_{Kahn} has m -torsion (Lemma 2.1), we can assume $\rho_{\text{Kahn}, A, k}([\zeta])$ to have values in $H_m^4(k)$.

Let k be the prime field of characteristic p and set $k' = k(\zeta)$ for $\zeta \in \bar{k}$ a primitive n^2 -th root of unity. Take $T = (t_1, t_2)_n \otimes (t_3, t_4)_n$ over $F = k'(t_1, t_2, t_3, t_4)$. We prove the formula for T . The proof ends by specialising to A as in the proof of Proposition 4.3.

Let $B : k\text{-fields} \rightarrow \mathfrak{Sets}$ be the functor attaching to a field extension F of k the Galois cohomology group $H^1(F, \mu_m)^4$ and H associating $H^4(F, \mu_m^{\otimes 4})$ with F . Now ρ_{Kahn} induces a cohomological invariant of B into H . Indeed, using the isomorphism $H^1(F, \mu_n) \cong F^\times / (F^\times)^n$, we associate with any four representants $a, b, c, d \in F^\times$ of classes in $H^1(F, \mu_m)$ the value $\rho_{\text{Kahn}, A, F}([\zeta]) \in H_m^4(F) \cong H^4(F, \mu_m^{\otimes 4}) \cong K_4^M(F)/m$ (for $A = (a, b)_n \otimes (c, d)_n$).

Using a full description of all possible invariants of B into H of [Gar, Prop. 2.1 & §3.1] and [GMS, Ex. 16.5], we find that $\rho_{\text{Kahn}, T, F}([\zeta])$ can be written in $K_4(F)/m$ as sum of pure symbols of the form $\lambda \{z_1, z_2, z_3, z_4\}$ where λ is an integer and each z_i is either a t_j either an element of k . We prove that only $\{t_1, t_2, t_3, t_4\}$ occurs. By specializing t_1 to 1, we obtain $T_1 = (1, t_2)_n \otimes (t_3, t_4)_n$ from T . But then $\mathbf{SK}_1(T_1) = 0$ by Wang's theorem so that $\rho_{\text{Kahn}, T_1, F}([\zeta]) = 0$. This induces that for all (non-trivial) pure symbols $\{z_1, z_2, z_3, z_4\}$ appearing in $\rho_{\text{Kahn}, T, F}([\zeta])$ one of the z_i has to equal t_1 (as the other ones are zero by the specialisation above). Three other specialisations give the result. ■

Remark 4.10 – In wild characteristics (i.e. when $p | n$), a formula as above does not make sense as there are no non-trivial p^2 -th roots of unity. So similar as in §3.4, we cannot generalise this formula to wild invariants by means of a lift.

(b) *Non-triviality of factor* – We prove the non-triviality of the factor appearing in Proposition 4.8. This uses the non-triviality of ρ_{Kahn} for Platonov's examples (Theorem 4.5). First we recall some notions related to tori. See [CTS] as a reference for more details.

Denote for a finite separable field extension K of k by $R_{K/k}(\mathbb{G}_m)$ the torus obtained by Weil restriction of scalars from K to k . Denote furthermore the kernel of the multiplication map $R_{K/k}(\mathbb{G}_m) \rightarrow \mathbb{G}_{m, k}$ by $R_{K/k}^1(\mathbb{G}_m)$ and the cokernel of the injection $\mathbb{G}_{m, k} \rightarrow R_{K/k}(\mathbb{G}_m)$ by $R_{K/k}(\mathbb{G}_m)/\mathbb{G}_m$. Furthermore for any k -torus T , we denote

by $T(k)/R$ the R -equivalence classes of $T(k)$. The dual $\hat{T}$ of a k -torus T is the character group $\text{Hom}(T, \mathbb{G}_m)$. The dual of $R_{K/k}(\mathbb{G}_m)$ is clearly the free abelian group $\mathbb{Z}[\Gamma]$ for $\Gamma = \text{Gal}(K/k)$. The dual of $R_{K/k}^1(\mathbb{G}_m)$ is then J_Γ , the cokernel of the norm:

$$\mathbb{Z} \rightarrow \mathbb{Z}[\Gamma] : a \mapsto \sum_{\gamma_i \in \Gamma} a\gamma_i.$$

The dual of $R_{K/k}(\mathbb{G}_m)/\mathbb{G}_m$ is the kernel I_Γ of the augmentation map:

$$\mathbb{Z}[\Gamma] \rightarrow \mathbb{Z} : \sum_{\gamma_i \in \Gamma} n_i \gamma_i \mapsto \sum_{\gamma_i \in \Gamma} n_i.$$

Recall that a k -torus F is called *flabby* (*flasque*) if $\hat{F}$ is a flabby Γ_k -module, i.e. $\text{Ext}^1(\hat{F}, P) = 0$ for any permutation Γ_k module P (for equivalent definitions see *ibid.*, Lem. 1). A flasque resolution of a k -torus T is an exact sequence of k -tori

$$0 \rightarrow S \rightarrow E \rightarrow T \rightarrow 0$$

with E quasi-trivial (i.e. $\hat{E}$ is a permutation module) and S flabby. This always exists and if T is split by a field extension K , then E and S can also be chosen to be split by K .

Theorem 4.11. *Let k be a p -adic field containing a n^3 -th primitive root unity. Suppose $A = (a, t_1)_n \otimes (c, t_2)_n$ is a division $k((t_1))((t_2))$ -algebra, then*

$$\rho_{Kahn,A,k}([\zeta]) = \varphi [\lambda h_n^4(\{a, t_1, c, t_2\})] \in H_{n^2}^4(k)$$

for ζ an n^2 -th primitive root of unity and an integer $\lambda \not\equiv 0 \pmod{\overline{n^2}}$ (and φ as in Proposition 4.8). A fortiori, $j(0, n) \not\equiv 0 \pmod{\overline{n^2}}$ for any n .

Proof. We know by Theorem 4.5 that $\rho_{Kahn,A} : \mathbf{SK}_1(A)(k) \rightarrow H_{n^2}^4(k)$ is not trivial and moreover $\mathbf{SK}_1(A)(k) = \mathbb{Z}/n$ and $H_{n^2}^4(k) \cong \mathbb{Z}/n^2$. We prove that the image of $\mu_{n^2}(k) \cong \mathbb{Z}/n^2$ inside $\mathbf{SK}_1(A)(k)$ is all of $\mathbf{SK}_1(A)(k)$. In that case, $\rho_{Kahn,A}([\zeta])$ is not trivial in $H_{n^2}^4(k)$ (and in $H_{n^2}^4(k) \cong \mathbb{Z}/\overline{n^2}$) so that $j(0, n) \not\equiv 0 \pmod{\overline{n^2}}$.

To prove the statement, let $L = k(\sqrt[n]{a}, \sqrt[n]{b})$ and $\Gamma = \text{Gal}(L/k) \cong \mathbb{Z}/n \times \mathbb{Z}/n$. Then by taking residues on $k((t_1))((t_2))$ with respect to t_1 and t_2 , Platonov proves $\mathbf{SK}_1(A)(k) \cong \hat{H}^{-1}(\Gamma, L^\times)$ where the cohomology group is a Tate cohomology group (see e.g. [Wei1, Def. 6.2.4]) - also use [Pla, Thms. 4.17 & 5.7] and [Wad, (6.15)]). On the other hand, $\hat{H}^{-1}(\Gamma, L^\times) = T(k)/R$ for $T = R_{L/k}^1(\mathbb{G}_m)$ [CTS, Prop. 15]. The resulting isomorphism $\mathbf{SK}_1(A)(k) \cong T(k)/R$ is a specialisation morphism (in t_1 and t_2) [Wad, (6.9) & (6.10)] so that the composite $\mu_{n^2}(k) \rightarrow \mathbf{SK}_1(A)(k) \cong T(k)/R$ is the canonical morphism $\mu_{n^2}(k) \rightarrow T(k)/R$. It suffices to prove that the latter is surjective.

First take a flabby resolution $1 \rightarrow S \rightarrow E \rightarrow T \rightarrow 1$ of L -split tori, then $H^1(k, S) = T(k)/R$ (*loc. cit.*, Thm. 2). The evaluation morphism $S \times \hat{S} \rightarrow \mathbb{G}_m$ induces a perfect pairing [Nak, Tat]:

$$H^1(k, S) \times H^1(k, \hat{S}) \rightarrow H^2(k, \mathbb{G}_m) \cong \mathbb{Q}/\mathbb{Z}.$$

Moreover $H^1(k, S) \cong H^1(\Gamma, S(L))$ as this follows from the inflation-restriction exact sequence [GS, 3.3.14] and $H^1(L, S) = 0$. The pairing above can be modified to a pairing

$$H^1(\Gamma, S(L)) \times H^1(\Gamma, \hat{S}(L)) \rightarrow \text{Br}(L/k) \cong \mathbb{Z}/n^2\mathbb{Z}.$$

Note that $\mu_{n^2}(k) \subset T$ so that we get a dual map $\hat{T} \rightarrow \mathbb{Z}/n^2\mathbb{Z}$. Using the flabby resolution and the pairing $T(k) \times \hat{T}(L) \rightarrow L^\times$, we get the following commutative diagram of pairings:

$$\begin{array}{ccccc}
H^1(k, S) & \times & H^1(k, \hat{S}) & \longrightarrow & H^2(k, \mathbb{G}_m) \cong \mathbb{Q}/\mathbb{Z} \\
\cong \uparrow & & \downarrow \cong & & \uparrow \\
H^1(\Gamma, S(L)) & \times & H^1(\Gamma, \hat{S}(L)) & \longrightarrow & \text{Br}(L/k) \\
\uparrow & & \downarrow & & \parallel \\
T(k) & \times & H^2(\Gamma, \hat{T}(L)) & \longrightarrow & \text{Br}(L/k) \\
\uparrow & & \downarrow & & \parallel \\
\mu_{n^2}(k) & \times & H^2(\Gamma, \mathbb{Z}/n^2) & \longrightarrow & \text{Br}(L/k).
\end{array}$$

The bottom pairing is perfect as $\mu_{n^2}(k) \cong \mathbb{Z}/n^2$; note that the bottom square comes from the compatibility of the pairings

$$\begin{array}{ccc}
T(k) & \times & \hat{T}(L) \longrightarrow L^\times \\
\uparrow & & \downarrow \\
\mu_{n^2}(k) & \times & \mathbb{Z}/n^2 \longrightarrow L^\times.
\end{array}$$

As $H^1(k, S) = T(k)/R \cong \mathbb{Z}/n$, to prove the surjectivity of $\mu_{n^2} \rightarrow T(k)/R$ it suffices to prove the injectivity of $H^1(k, \hat{S}) \rightarrow H^2(\Gamma, \mathbb{Z}/n^2)$. Since $H^1(\Gamma, \hat{E}(L)) = 0$, this comes down to proving the injectivity of $H^2(\Gamma, \hat{T}) \rightarrow H^2(\Gamma, \mathbb{Z}/n^2)$. This morphism fits into an exact sequence

$$H^2(\Gamma, I_\Gamma) \rightarrow H^2(\Gamma, \hat{T}) \rightarrow H^2(\Gamma, \mathbb{Z}/n^2)$$

because of the exact sequence of group functors

$$0 \rightarrow \mu_{n^2} \rightarrow T \rightarrow R_{L/k}(\mathbb{G}_m)/\mathbb{G}_m \rightarrow 0.$$

Clearly $T \rightarrow R_{L/k}(\mathbb{G}_m)/\mathbb{G}_m$ factors through $R_{L/k}(\mathbb{G}_m)$, so that $H^2(\Gamma, I_\Gamma) \rightarrow H^2(\Gamma, \hat{T})$ factors through $H^2(\Gamma, \mathbb{Z}[\Gamma])$ which is trivial by Shapiro's Lemma. This proves the desired injectivity. $\blacksquare$

Remark 4.12 – Note that the proof also defines an invariant of the torus T with values inside $H_{n^2}^4(k)$.

From this we get the following corollary.

Corollary 4.13. *Let k be a field containing an l^2 -th primitive root of unity (for $l \neq \text{char}(k)$ any prime) and let $A = (a, b)_l \otimes (c, d)_l$ be a product of two symbol algebras. If $\{a, b, c, d\} \neq 0 \in K_4^M(k)/l$, then $\mathbf{SK}_1(A) \neq 0$.*

Proof. For a field k of characteristic 0, the corollary follows from the previous theorem.

Let k be a field of $\text{char}(k) = p > 0$ and let $l \neq p$ be a prime and assume k to contain an l^2 -th primitive root $\zeta \in \bar{k}$. Take any product of two symbol k -algebras $A = (a, b)_l \otimes (c, d)_l$ for $a, b, c, d \in k^\times$. Let R be a p -ring with residue field k and fraction field K . Then

A lifts to the central simple K -algebra $B = (\tilde{a}, \tilde{b})_l \otimes (\tilde{c}, \tilde{d})_l$ where $\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}$ are lifts from $a, b, c, d \in R$. Under the injection $H_{l^2}^4(k) \rightarrow H_{l^2}^4(K)$ induced by (2.5), $\varphi[h_{l,k}^4(\{a, b, c, d\})]$ is sent to $\varphi[h_{l,K}^4(\{\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}\})]$ (with an abuse of notation for φ from Proposition 4.8). This follows from a same splitting for Milnor's K-Theory [Mil, Lem. 2.6].

As $\rho_{\text{Kahn}, B, K}([\tilde{\zeta}]) = \varphi[j(0, l) h_{l,K}^4(\{\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}\})]$ (for a lift $\tilde{\zeta} \in \mu_{l^2}(K)$ of ζ), we find from the construction in Theorem 2.4 that $\tilde{\rho}_{\text{Kahn}, A, k}([\zeta]) = \varphi[j(0, l) h_{l,k}^4(\{a, b, c, d\})]$. On the other hand, as Kahn's invariant generates the invariant group (§2.4 (a)), there is an integer d such that $\tilde{\rho}_{\text{Kahn}, A} = d \rho_{\text{Kahn}, A}$. From this the result follows. ■

By the proof, it even suffices to prove the non-triviality of the symbols in characteristic zero to obtain the non-triviality of the symbols in moderate positive characteristic. (Use a p -ring and the splitting in Milnor's K-Theory (loc. cit.)).

References

- [AG] Maurice Auslander and Oscar Goldman. The Brauer group of a commutative ring. *Trans. Amer. Math. Soc.*, 97:367–409, 1960.
- [Alb] Adrian Albert. Normal division algebras of degree four over an algebraic field. *Trans. Amer. Math. Soc.*, 34(2):363–372, 1932.
- [Bae] Ricardo Baeza. *Quadratic forms over semilocal rings*. Lecture Notes in Mathematics, Vol. 655. Springer-Verlag, Berlin, 1978.
- [BK] Spencer Bloch and Kazuya Kato. p -adic étale cohomology. *Publ. Math. Inst. Hautes Études Sci.*, (63):107–152, 1986.
- [Coh] Irvin Cohen. On the structure and ideal theory of complete local rings. *Trans. Amer. Math. Soc.*, 59:54–106, 1946.
- [CTS] Jean-Louis Colliot-Thélène and Jean-Jacques Sansuc. La R -équivalence sur les tores. *Ann. Sci. École Norm. Sup. (4)*, 10(2):175–229, 1977.
- [Dra] Peter Draxl. *Skew Fields*, volume 81 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1983.
- [Gar] Skip Garibaldi. Cohomological invariants: exceptional groups and spin groups. *Mem. Amer. Math. Soc.*, 200(937):xii+81, 2009. With an appendix by Detlev W. Hoffmann.
- [GMS] Skip Garibaldi, Alexander Merkurjev, and Jean-Pierre Serre. *Cohomological invariants in Galois cohomology*, volume 28 of *University Lecture Series*. Amer. Math. Soc., 2003.
- [Gro1] Alexander Grothendieck. *Éléments de Géométrie Algébrique IV, Étude locale des schémas et des morphismes de schémas, Première Partie*, volume 20 of *Publ. Math. Inst. Hautes Études Sci.* Bures-sur-Yvette, 1964.
- [Gro2] Alexander Grothendieck. Le groupe de Brauer : I. Algèbres d'Azumaya et interprétations diverses. *Séminaire Bourbaki*, 9:199–219, 1964–1966. Exposé No. 290.
- [GS] Philippe Gille and Tamás Szamuely. *Central Simple Algebras and Galois Cohomology*, volume 101 of *Cambridge studies in advanced mathematics*. Cambridge University Press, Cambridge, 2006.
- [Izh] Oleg Izhboldin. On the cohomology groups of the field of rational functions. In *Mathematics in St. Petersburg*, volume 174 of *Amer. Math. Soc. Transl. Ser. 2*, pages 21–44. Amer. Math. Soc., Providence, RI, 1996.
- [Kah1] Bruno Kahn. *Formes quadratiques sur un corps*, volume 15 of *Cours Spécialisés*. Société Mathématique de France, 2008.
- [Kah2] Bruno Kahn. Cohomological approaches to SK_1 and SK_2 of central simple algebras. *Preprint*, 2009.
- [Kat1] Kazuya Kato. Galois cohomology of complete discrete valuation fields. In *Algebraic K-Theory*, volume 967 of *Lecture notes in mathematics*, pages 215–238, Berlin, 1982.
- [Kat2] Kazuya Kato. Symmetric bilinear forms, quadratic forms and Milnor K -theory in characteristic two. *Invent. Math.*, 66(3):493–510, 1982.

- [KMRT] Max-Albert Knus, Alexander Merkurjev, Markus Rost, and Jean-Pierre Tignol. *The book of involutions*, volume 44 of *Amer. Math. Soc. Colloq. Publ.* 1998.
- [Knu] Max-Albert Knus. *Quadratic and Hermitian forms over rings*, volume 294 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin, 1991.
- [Mat] Hideyuki Matsumura. *Commutative ring theory*, volume 8 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1986. Translated from the Japanese by M. Reid.
- [Mer1] Alexander Merkurjev. K -theory of simple algebras. In *K-theory and algebraic geometry: connections with quadratic forms and division algebras (Santa Barbara, CA, 1992)*, volume 58 of *Proc. Sympos. Pure Math.*, pages 65–83. Amer. Math. Soc., Providence, RI, 1995.
- [Mer2] Alexander Merkurjev. Invariants of algebraic groups. *J. reine angew. Math.*, 508:127–156, 1999.
- [Mer3] Alexander Merkurjev. The group SK_1 for simple algebras. *K-Theory*, 37(3):311–319, 2006.
- [Mil] John Milnor. Algebraic K -theory and quadratic forms. *Invent. Math.*, 9:318–344, 1969/1970.
- [Nak] Tadasi Nakayama. Cohomology of class field theory and tensor product modules. I. *Ann. of Math. (2)*, 65:255–267, 1957.
- [NM] Tadasi Nakayama and Yozô Matsushima. Über die multiplikative Gruppe einer p -adischen Divisionsalgebra. *Proc. Imp. Acad. Tokyo*, 19:622–628, 1943.
- [OVV] Dmitri Orlov, Alexander Vishik, and Vladimir Voevodsky. An exact sequence for $K_*^M/2$ with applications to quadratic forms. *Ann. of Math.*, 165(1):1–13, 2007.
- [Pla] Vladimir Platonov. The Tannaka-Artin problem and reduced K -theory. *Math. USSR Izv.*, 10(2):211–243, 1976. English translation.
- [Ros1] Markus Rost. Chow Groups with Coefficients. *Doc. Math. J. DMV*, 1:319–393, 1996.
- [Ros2] Markus Rost. The basic correspondence of a splitting variety. 1998. Notes downloadable from his website.
- [RTY] Ulf Rehmann, Sergey Tikhonov, and Vyacheslav Yanchevskiĭ. Symbols and cyclicity of algebras after a scalar extension. *Fundam. Prikl. Mat.*, 14(6):193–209, 2008.
- [Ser] Jean-Pierre Serre. *Galois Cohomology*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2002.
- [SGA] *Schémas en groupes. III: Structure des schémas en groupes réductifs*. Séminaire de Géométrie Algébrique du Bois Marie 1962/64 (SGA 3). Dirigé par M. Demazure et A. Grothendieck. Lecture Notes in Mathematics, Vol. 153. Springer-Verlag, Berlin, 1962/1964.
- [Sus1] Andrei Suslin. SK_1 of division algebras and Galois cohomology. In *Algebraic K-theory*, volume 4 of *Adv. Soviet Math.*, pages 75–99. Amer. Math. Soc., Providence, RI, 1991.
- [Sus2] Andrei Suslin. SK_1 of division algebras and Galois cohomology revisited. In *Proceedings of the St. Petersburg Mathematical Society. Vol. XII*, volume 219 of *Amer. Math. Soc. Transl. Ser. 2*, pages 125–147, Providence, RI, 2006. Amer. Math. Soc.
- [Tat] John Tate. The cohomology groups of tori in finite Galois extensions of number fields. *Nagoya Math. J.*, 27:709–719, 1966.
- [Voe] Vladimir Voevodsky. On Motivic Cohomology with $\mathbb{Z}/l$ coefficients. *Preprint*, 2009.
- [Wad] Adrian Wadsworth. Valuation theory on finite dimensional division algebras. In *Valuation theory and its applications, Vol. I (Saskatoon, SK, 1999)*, volume 32 of *Fields Inst. Commun.*, pages 385–449. Amer. Math. Soc., Providence, RI, 2002.
- [Wan] Shianghaw Wang. On the commutator group of a simple algebra. *Amer. J. Math.*, 72:323–334, 1950.
- [Wei1] Charles Weibel. *An introduction to homological algebra*, volume 38 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1997.
- [Wei2] Charles Weibel. The norm residue isomorphism theorem. *J. Topol.*, 2(2):346–372, 2009.
- [Wit] Ernst Witt. Zyklische Körper und Algebren der Charakteristic p vom Grad p^n . *J. reine angew. Math.*, 176:126–140, 1937.
- [Wou] Tim Wouters. L’invariant de Suslin en caractéristique positive. *Preprint*, 2009.